

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 January 2004 (08.01.2004)

PCT

(10) International Publication Number
WO 2004/002986 A2

(51) International Patent Classification⁷: **C07D 471/04**,
487/04, 473/00, 405/08, A61K 31/41, 31/435, 31/52,
A61P 25/00, 9/00, 3/00

(21) International Application Number:
PCT/JP2003/008161

(22) International Filing Date: 26 June 2003 (26.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2002-190978 28 June 2002 (28.06.2002) JP

(71) Applicant (for all designated States except US): **BANYU PHARMACEUTICAL CO., LTD.** [JP/JP]; 2-3, Nihonbashi Honcho 2-Chome, Chou-ku, Tokyo 103-8416 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OTAKE, Norikazu** [JP/JP]; c/o BANYU PHARMACEUTICAL CO., LTD. Tsukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki 300-2611 (JP). **MORIYA, Minoru** [JP/JP]; c/o BANYU PHARMACEUTICAL CO., LTD. Tsukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki 300-2611 (JP). **OGINO, Yoshio** [JP/JP]; c/o BANYU PHARMACEUTICAL CO., LTD., Tsukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki 300-2611 (JP). **MATSUDA, Kenji** [JP/JP]; c/o BANYU PHARMACEUTICAL CO.,

LTD., 2-3, Nihonbashi Honcho 2-chome, Chuo-ku, TOKYO 103-8416 (JP). **NAGAE, Yoshikazu** [JP/JP]; c/o BANYU PHARMACEUTICAL CO., LTD., Tsukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki 300-2611 (JP). **KANATANI, Akio** [JP/JP]; c/o BANYU PHARMACEUTICAL CO., LTD., Tsukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki 300-2611 (JP). **FUKAMI, Takehiro** [JP/JP]; c/o BANYU PHARMACEUTICAL CO., LTD., Tsukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki 300-2611 (JP).


(74) Agent: **IWATANI, Ryo**; Sakurabashi Chiyoda Build., 5F, 1-27, Dojima 2-chome, Kita-ku, Osaka-shi, Osaka 530-0003 (JP).

(81) Designated States (national): AE, AG, AL, AM, AU, AZ, BA, BB, BR, BY, BZ, CA, CN, CO, CR, CU, DM, DZ, EC, GD, GH, HR, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MA, MD, MG, MK, MN, MX, NI, NO, NZ, OM, PG, PH, PL, RU, SC, SG, SY, TJ, TM, TN, TT, UA, US, UZ, VC, VN, YU, ZA.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: NOVEL BENZIMIDAZOLE DERIVATIVES



(1)

(57) Abstract: A compound of the formula (I): (wherein A, B, C and D are independently nitrogen or optionally substituted methine; E is nitrogen, methine or hydroxy substituted methine; n is 0 or 1; T, U, V and W are independently nitrogen or optionally substituted methine; X is N(SO₂R⁴), -N(COR⁵) or -CO-; Y is -C(R⁶)(R⁷), -O- or -N(R⁸), provided that the compound (I) when E is nitrogen, n is 0, X is -CO-, and Y is -O- is excluded) is useful

as an agent for the treatment of various diseases related to NPY, for example cardiovascular disorders such as angina, acute or congestive heart failure, myocardial infarction, hypertension, nephropathy, electrolyte abnormality, vasospasm, arteriosclerosis, etc., central nervous system disorders such as bulimia, depression, anxiety, seizure, epilepsy, dementia, pain, alcoholism, drug withdrawal, circadian rhythm disorders, schizophrenia, memory impairment, sleep disorders, cognitive impairment, etc., metabolic diseases such as obesity, diabetes, hormone abnormality, hypercholesterolemia, hyperlipidemia, gout, fatty liver, etc., genital or reproductive disorders such as infertility, preterm labor, sexual dysfunction, etc., gastro-intestinal disorders, respiratory disorder, inflammatory diseases or glaucoma, and the like, also for example, atherosclerosis, hypogonadism, hyperandrogenism, polycystic ovary syndrome, hirsutism, gastro-intestinal motility disorder, obesity-related gastro-esophageal reflux, obesity hypoventilation (Pickwickian syndrome), sleep apnea, inflammation, systemic inflammation of the vasculature, osteoarthritis, insulin resistance, bronchoconstriction, alcohol preference, metabolic syndrome, Alzheimer's disease, cardiac hypertrophy, left ventricular hypertrophy, hypertriglyceridemia, low HDL cholesterol, cardiovascular disorders such as coronary heart disease (CHD), cerebrovascular disease, stroke, peripheral vascular disease, sudden death, gallbladder diseases, cancer (breast, endometrial, colon), breathlessness, hyperuricemia, impaired fertility, low back pain, and increased anesthetic risk.

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AU, AZ, BA, BB, BR, BY, BZ, CA, CN, CO, CR, CU, DM, DZ, EC, GD, GH, HR, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MA, MD, MG, MK, MN, MX, NI, NO, NZ, OM, PG, PH, PL, RU, SC, SG, SY, TJ, TM, TN, TT, UA, UZ, VC, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,

GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- of inventorship (Rule 4.17(iv)) for US only

Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

DESCRIPTION

NOVEL BENZIMIDAZOLE DERIVATIVES

5 Technical Field

The present invention is useful in medical fields. In more detail, novel benzimidazole derivatives of the present invention have an effect as neuropeptide Y receptor antagonists and are useful as agents for the treatment of various kinds of
10 cardiovascular disorders, central nervous system disorders, metabolic diseases and the like.

Background Art

Neuropeptide Y (hereinafter referred to as NPY), a
15 peptide consisting of 36 amino acids, was first isolated from porcine brain by Tatemoto et al in 1982 (NATURE, vol. 296, p. 659(1982)). NPY is widely distributed in central nervous system and peripheral nervous system, and plays various roles as one of the most abundant peptides in the nervous system. That
20 is, NPY acts as an orexigenic substance in the central nervous system and markedly promotes fat accumulation via the mediation of secretion of various hormones or the action of the nervous system. It is known that continuous intracerebroventricular administration of NPY induces obesity and insulin resistance
25 due to these actions (INTERNATIONAL JOURNAL OF OBESITY, vol.19, p.517(1995); Endocrinology, vol.133, p.1753(1993)). It is also known that NPY has central actions such as depression, anxiety, schizophrenia, pain, dementia, circadian rhythm control and the like (DRUGS, vol.52, p.371(1996); THE JOURNAL

OF NEUROSCIENCE, vol.18, p.3014(1998)). Furthermore, in the periphery, NPY coexists with norepinephrine in sympathetic-nerve terminals and is related to the tonicity of the sympathetic nervous system. It is known that peripheral
5 administration of NPY causes vasoconstriction and enhances the activities of other vasoconstrictive substances such as norepinephrine (BRITISH JOURNAL OF PHARMACOLOGY, vol.95, p.419(1988)). It is also reported that NPY could participate in the development of cardiac hypertrophy as a result of the
10 sympathetic stimulation (PROCEEDING NATIONAL ACADEMIC SCIENCE USA, vol.97, p.1595(2000)) .

On the other hand, it is reported that NPY is also involved in the secretory function of sexual hormones and growth hormone, sexual behavior and reproductive function, gastro-intestinal
15 motility, bronchoconstriction, inflammation and alcohol preference (LIFE SCIENCE, vol.55, p.551(1994); THE JOURNAL OF ALLERGY AND IMMUNOLOGY, vol.101, p.S345(1998); NATURE, vol.396, p.366(1998)).

NPY has a variety of pharmacological effects resulting from NPY binding to the NPY receptors, to some of which other
20 NPY related peptides including peptide YY and pancreatic polypeptide also bind. It is known that these pharmacological effects of NPY are mediated by the action of at least five receptors with or without synergistic interactions (TRENDS IN
25 NEUROSCIENCE, vol.20, p.294(1997)).

It is reported that the central effects mediated by NPY Y1 receptor include remarkable orexigenic effect (ENDOCRINOLOGY, vol.137, p.3177(1996); ENDOCRINOLOGY, vol.141, p.1011(2000)). Further, NPY Y1 receptor is reported to be

involved in anxiety and pain (NATURE, vol.259, p.528(1993);
BRAIN RESEARCH, vol.859, p.361(2000). In addition, the pressor
effect mediated by the strong vasoconstrictor action in the
periphery is also reported (FEBS LETTERS, vol.362, p.192(1995);
5 NATURE MEDICINE, vol.4, p.722(1998)).

It is known that the effects mediated by NPY Y2 receptor
include an inhibitory effect on the release of various
neurotransmitters in the sympathetic nerve endings (BRITISH
JOURNAL OF PHARMACOLOGY, vol.102, p.41(1991); SYNAPSE, vol.2,
10 p.299(1988)). In periphery, NPY Y2 causes constriction of blood
vessel or vas deferens directly or via the control of release
of various neurotransmitters (THE JOURNAL OF PHARMACOLOGY AND
EXPERIMENTAL THERAPEUTICS, vol.261, p.863(1992); BRITISH
JOURNAL OF PHARMACOLOGY, vol.100, p.190(1990)). Inhibition of
15 lipolysis in adipose tissues is also known (ENDOCRINOLOGY,
vol.131, p.1970(1992)). Further, inhibition of ion secretion
in the gastro-intestinal tract is reported (BRITISH JOURNAL OF
PHARMACOLOGY, vol.101, p.247(1990)). On the other hand, the
effect on the central nervous system functions such as memory,
20 anxiety and the like are also known. (BRAIN RESEARCH, vol.503,
p.73(1989); PEPTIDES, vol.19, p.359(1998)).

It is reported that NPY Y3 receptor exists mainly in
brainstem and heart and is related to the regulation of blood
pressure and heart rate (THE JOURNAL OF PHARMACOLOGY AND
25 EXPERIMENTAL THERAPEUTICS, vol.258, p.633(1991); PEPTIDES,
vol.11, p.545(1990)). It is also known that NPY Y3 is involved
in the control of catecholamine secretion in adrenal gland (THE
JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, vol.244,
p.468(1988); LIFE SCIENCE, vol.50, p.PL7(1992)).

NPY Y4 receptor has high affinity for pancreatic polypeptide in particular. As for the pharmacological effects of NPY Y4, inhibition of pancreatic exocrine secretion and gastro-intestinal motility is reported (GASTROENTEROLOGY, 5 vol.85, p.1411(1983)). Further, it is reported that NPY enhances the secretion of sexual hormones in the central nervous system (ENDOCRINOLOGY, vol.140, p.5171(1999)).

As for the effects mediated by NPY Y5 receptor, fat accumulation effects including orexigenic effect are prominent 10 (NATURE, vol. 382, p.168(1996); AMERICAN JOURNAL OF PHYSIOLOGY, vol.277, p.R1428(1999)). It is also reported that the NPY Y5 receptor mediates some CNS effects, such as seizure and epilepsy, or pain and morphine withdrawal symptoms, and the control of circadian rhythm (NATURE MEDICINE, vol.3, 15 p.761(1997); PROCEEDING NATIONAL ACADEMIC SCIENCE USA, vol.96, p.13518(1999); THE JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, vol.284, p.633(1998); THE JOURNAL OF NEUROSCIENCE, vol.21, p.5367(2001). In addition, diuresis effect and hypoglycemic effect in the periphery are reported 20 (BRITISH JOURNAL OF PHARMACOLOGY, vol. 120, p.1335(1998); ENDOCRINOLOGY, vol.139, p.3018(1998)). NPY is also reported to enhance cardiac hypertrophy as a result of the sympathetic accentuation (PROCEEDING NATIONAL ACADEMIC SCIENCE USA, vol.97, p.1595 (2000)).

25 The effects of NPY are expressed when NPY binds to the NPY receptors in the central or peripheral nervous system. Therefore, the action of NPY can be prevented by blocking its binding to NPY receptors. For this reason, it is expected that substances antagonize NPY binding to NPY receptors may be useful

for the prophylaxis or treatment of various diseases related to NPY, for example cardiovascular disorders such as angina, acute or congestive heart failure, myocardial infarction, hypertension, nephropathy, electrolyte abnormality, vasospasm, atherosclerosis, etc., central nervous system disorders such as bulimia, depression, anxiety, seizure, epilepsy, dementia, pain, alcoholism, drug withdrawal, circadian rhythm disorders, schizophrenia, memory impairment, sleep disorders, cognitive impairment, etc., metabolic diseases such as obesity, diabetes, hormone abnormality, gout, fatty liver, etc., genital or reproductive disorders such as infertility, preterm labor, sexual dysfunction, etc., gastro-intestinal disorders, respiratory disorder, inflammatory diseases or glaucoma, and the like. (TRENDS IN PHARMACOLOGICAL SCIENCE, vol.15, p.153(1994); LIFE SCIENCE, vol.55, p.551(1994); DRUGS, vol.52, p.371(1996); THE JOURNAL OF ALLERGY AND IMMUNOLOGY, vol.101, p.S345(1998); NATURE, vol.396, p.366(1998); THE JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, vol.284, p.633(1998); TRENDS IN PHARMACOLOGICAL SCIENCE, vol.20, p.104(1999); PROCEEDING NATIONAL ACADEMIC SCIENCE USA, vol.97, p.1595(2000); THE JOURNAL OF NEUROSCIENCE, vol. 21, p.5367(2001); PHARMACOLOGY & THERAPEUTICS, vol.65, p.397(1995)).

It was recently found that, as a result of the study by the present inventors, a certain NPY receptor antagonist is useful for the prophylaxis or treatment of hypercholesterolemia, hyperlipidemia and arteriosclerosis (International application publication WO99/27965).

International application publication WO00/27845 and


WO01/14376 disclose a variety of carboxamide derivatives, and mention that said derivatives have excellent NPY receptor antagonistic activities. International application publication WO02/48152 discloses a variety of
 5 spiro(isobenzofuran-1,4'-piperidine)-3-on derivatives, and mentions that said derivatives have an effect to regulate the NPY binding to NPY5 receptors. However, none of the above international publications describes the compounds of the present invention.

10

Disclosure of Invention

The object of the present invention is to provide novel medicines which exhibit NPY antagonistic activities.

The present inventors have discovered that the compounds
 15 of the general formula (I):



(wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower
 20 alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R¹)R² and -Q¹-Ar¹, and at least one of A, B, C and D is said methine group;

Ar¹ is aryl or heteroaryl, any of which is optionally
 25 substituted by a substituent selected from the group consisting

of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxy carbonyl,
5 lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower
10 alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine (namely methine substituted by hydroxy);

n is 0 or 1;

Q^1 and Q^2 are independently a single bond, oxygen, carbonyl
15 or $-N(R^3)-$;

R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

20 R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;

R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;

25 T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group;

X is $-\text{N}(\text{SO}_2\text{R}^4)-$, $-\text{N}(\text{COR}^5)-$ or $-\text{CO}-$;

Y is $-\text{C}(\text{R}^6)(\text{R}^7)-$, $-\text{O}-$ or $-\text{N}(\text{R}^8)-$, provided that the compound (I) when E is nitrogen, n is 0, X is $-\text{CO}-$ and Y is $-\text{O}-$ is excluded)

5 exhibit NPY antagonistic activities especially on NPY Y5 receptors and show excellent pharmacokinetics such as transport into brain or transport to cerebrospinal fluid, etc., thereby completed the present invention.

Compounds of the present invention (I) exhibit NPY
10 antagonistic effects especially on NPY Y5 receptors and show excellent pharmacokinetics such as transport into brain or transport to cerebrospinal fluid, etc. Also, the compound of the present invention (I) are useful as agents for the treatment of various diseases related to NPY, for example, cardiovascular
15 disorders such as angina, acute or congestive heart failure, myocardial infarction, hypertension, nephropathy, electrolyte abnormality, vasospasm, arteriosclerosis, etc., central nervous system disorders such as bulimia, depression, anxiety, seizure, epilepsy, dementia, pain, alcoholism, drug
20 withdrawal, circadian rhythm disorders, schizophrenia, memory impairment, sleep disorders, cognitive impairment, etc., metabolic diseases such as obesity, diabetes, hormone abnormality, hypercholesterolemia, hyperlipidemia, gout, fatty liver, etc., genital or reproductive disorders such as
25 infertility, preterm labor, sexual dysfunction, etc., gastro-intestinal disorders, respiratory disorder, inflammatory diseases or glaucoma, and the like, also for example, atherosclerosis, hypogonadism, hyperandrogenism, polycystic ovary syndrome, hirsutism, gastro-intestinal

motility disorder, obesity-related gastro-esophageal reflux, obesity hypoventilation (Pickwickian syndrome), sleep apnea, inflammation, systemic inflammation of the vasculature, osteoarthritis, insulin resistance, bronchoconstriction, alcohol preference, metabolic syndrome, Alzheimer's disease, cardiac hypertrophy, left ventricular hypertrophy, hypertriglyceridemia, low HDL cholesterol, cardiovascular disorders such as coronary heart disease (CHD), cerebrovascular disease, stroke, peripheral vascular disease, sudden death, gallbladder diseases, cancer (breast, endometrial, colon), breathlessness, hyperuricemia, impaired fertility, low back pain, or increased anesthetic risk, and the like.

The compounds of the present invention (I) are particularly useful as agents for the treatment of bulimia, obesity, diabetes and the like.

The present invention relates to the compounds represented by the general formula (I), or the salts thereof, and the production method and the use thereof.

The means of terms used in the present specification are defined and more detailed description of this invention is described in the following.

"Halogen" refers to fluorine, chlorine, bromine and iodine.

"Lower alkyl" refers to a straight- or branched-chain alkyl group of C1 to C6, and its examples are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, hexyl, isohexyl and the like.

"Halo-lower alkyl" refers to said lower alkyl substituted

with identically or differently one, two or more, preferably one to three said halogen at the substitutable, arbitrary position(s), and its examples are fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 1,2-difluoroethyl, chloromethyl, 2-chloroethyl, 1,2-dichloroethyl, bromomethyl, iodomethyl and the like.

"Lower alkoxy" refers to straight- or branched-chain alkoxy of C1 to C6 and its examples are methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, isobutoxy, tert-butoxy, pentyloxy, isopentyloxy, hexyloxy, isohexyloxy and the like.

"Halo-lower alkoxy" refers to said lower alkoxy substituted with identically or differently one, two or more, preferably one to three said halogen at substitutable, arbitrary position(s), and its examples are fluoromethoxy, difluoromethoxy, trifluoromethoxy, 2-fluoroethoxy, 1,2-difluoroethoxy, chloromethoxy, 2-chloroethoxy, 1,2-dichloroethoxy, bromomethoxy, iodomethoxy and the like.

"Lower alkoxy carbonyl" refers to an alkoxy carbonyl group containing said lower alkoxy, that is, an alkoxy carbonyl group of C2 to C7, and its examples are methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl and the like.

"Lower alkylsulfonyl" refers to a straight- or branched-chain alkylsulfonyl group of C1 to C6, and its examples are methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, sec-butylsulfonyl, isobutylsulfonyl, tert-butylsulfonyl, pentylsulfonyl, isopentylsulfonyl, hexylsulfonyl, isohexylsulfonyl and the

like.

"Lower alkylsulfonyloxy" refers to a straight- or branched-chain alkylsulfonyloxy group of C1 to C6, and its examples are methylsulfonyloxy, ethylsulfonyloxy, propylsulfonyloxy, isopropylsulfonyloxy, butylsulfonyloxy, sec-butylsulfonyloxy, isobutylsulfonyloxy, tert-butylsulfonyloxy, pentylsulfonyloxy, isopentylsulfonyloxy, hexylsulfonyloxy, isohexylsulfonyloxy and the like.

"Hydroxy-lower alkyl" refers to said lower alkyl substituted with one, two or more, preferably one or two hydroxy at substitutable, arbitrary position(s), and its examples are hydroxymethyl, 2-hydroxyethyl, 1-hydroxy-1-methylethyl, 1,2-dihydroxyethyl, 3-hydroxypropyl and the like.

"Cyclo-lower alkyl" refers to a cycloalkyl group of C3 to C6, and its examples are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

"Lower alkenyl" refers to a straight- or branched-chain alkenyl group of C2 to C6, and its examples are vinyl, 1-propenyl, 2-propenyl, isopropenyl, 3-butenyl, 2-butenyl, 1-butenyl, 1-methyl-2-propenyl, 1-methyl-1-propenyl, 1-ethyl-1-ethenyl, 2-methyl-2-propenyl, 2-methyl-1-propenyl, 3-methyl-2-butenyl, 4-pentenyl and the like.

"Lower alkylthio" refers to straight- or branched-chain alkylthio of C1 to C6, and its examples are methylthio, ethylthio, propylthio, isopropylthio, butylthio, sec-butylthio, isobutylthio, tert-butylthio, pentylthio, isopentylthio, hexylthio, isohexylthio and the like.

"Lower alkanoyl" refers to an alkanoyl group containing

said lower alkyl, that is, an alkanoyl group of C2 to C7, and its examples are acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl and the like.

"Lower alkanoylamino" refers to an amino group
5 mono-substituted with said lower alkanoyl, and its examples are acetilamino, propionylamino, butyrylamino, isobutyrylamino, valerylamino, isovalerylamino, pivaloylamino and the like.

"Aryl" refers to phenyl, naphthyl and the like.

"Heteroaryl" refers to 5- or 6-membered monocyclic
10 heteroaromatic group which contains one, two or more, preferably one to three hetero atom(s) identically or differently selected from the group consisting of oxygen, nitrogen and sulfur; or condensed cyclic heteroaromatic group, where said monocyclic heteroaromatic group is condensed with
15 said aryl group or condensed each other with the same or different said monocyclic heteroaromatic group, and its examples are pyrrolyl, furyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl, oxadiazolyl, 1,2,3-thiadiazolyl,
20 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, indolyl, benzofuranyl, benzothienyl, benzoimidazolyl, benzoxazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, indazolyl, purinyl, quinolyl, isoquinolyl,
25 phthalazinyl, naphthylidiny, quinoxaliny, quinazolinyl, cinnolinyl, pteridinyl, pyrido[3,2-b]pyridyl and the like.

"Lower alkylamino" refers to an amino group mono-substituted with said lower alkyl, and its examples are methylamino, ethylamino, propylamino, isopropylamino,

butylamino, sec-butylamino, tert-butylamino and the like.

"Di-lower alkylamino" refers to an amino group di-substituted with the same or different said lower alkyl, and its examples are dimethylamino, diethylamino, 5 ethylmethylamino, dipropylamino, methylpropylamino, diisopropylamino and the like.

"Lower alkylene which may be intervened by oxygen, sulfur or imino" refers to an alkylene group of C2 to C5, which is not intervened or intervened by one, two or more, preferably one 10 oxygen, sulfur or imino at optional and intervention capable position(s) of said alkylene chain, and its examples are ethylene, trimethylene, tetramethylene, pentamethylene, 2-oxatetramethylene, 2-oxapentamethylene, 3-oxapentamethylene, 2-thiatetramethylene, 15 2-thiapentamethylene, 3-thiapentamethylene, 2-azatetramethylene, 2-azapentamethylene, 3-azapentamethylene and the like.

"Aralkyl" refers to said lower alkyl substituted with one, two or more, preferably one said aryl at substitutable, 20 arbitrary position(s), and its examples are benzyl, 1-phenylethyl, phenethyl, 1-naphthylmethyl, 2-naphthylmethyl and the like.

The esters of compounds of formula (I) refer to, for example, the pharmaceutically acceptable, common esters of said 25 carboxyl group when the compound has a carboxyl group, and examples thereof are esters with lower alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, cyclopropyl, cyclobutyl, cyclopentyl), esters with aralkyl (e.g. benzyl, phenethyl), esters with lower

alkenyl (e.g. allyl, 2-butenyl), esters with lower-alkoxy-lower-alkyl (e.g. methoxymethyl, 2-methoxyethyl, 2-ethoxyethyl), esters with lower-alkanoyloxy-lower-alkyl (e.g. acetoxymethyl, pivaloyloxymethyl, 1-pivaloyloxyethyl),
5 esters with lower-alkoxycarbonyl-lower-alkyl (e.g. methoxycarbonylmethyl, isopropoxycarbonylmethyl), esters with carboxy-lower alkyl (e.g. carboxymethyl), esters with lower-alkoxycarbonyloxy-lower-alkyl (e.g. 1-(ethoxycarbonyloxy)ethyl,
10 1-(cyclohexyloxyloxy)ethyl), esters with carbamoyloxy-lower alkyl (e.g. carbamoyloxymethyl), esters with phthalidyl, esters with (5-substituted-2-oxo-1,3-dioxol-4-yl)methyl (e.g. (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl) and the like.

15 The salts of compounds of formula (I) refer to the pharmaceutically acceptable, common salts, and examples thereof are base addition salt to said carboxyl group when the compound has a carboxyl group, or acid addition salt to said amino or basic heterocyclyl when the compound has an amino or
20 basic heterocyclyl group and the like.

Said base addition salts include salts with alkali metals (e.g. sodium, potassium); salts with alkaline earth metals (e.g. calcium, magnesium); ammonium salts; salts with organic amines (e.g. trimethylamine, triethylamine, dicyclohexylamine,
25 ethanolamine, diethanolamine, triethanolamine, procaine, N,N'-dibenzylethylenediamine) and the like.

Said acid addition salts include salts with inorganic acids (e.g. hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid, perchloric acid), salts with organic acids

(e.g. maleic acid, fumaric acid, tartaric acid, citric acid, ascorbic acid, trifluoroacetic acid), salts with sulfonic acids (e.g. methanesulfonic acid, isethionic acid, benzenesulfonic acid, p-toluenesulfonic acid) and the like.

5 An N-oxide derivative of the compound represented by the formula (I) means a compound of which any one or more than one nitrogen atoms present in the compound of the formula (I) is or are oxidized to form an N-oxide or N-oxides, and such an N-oxide derivative includes, for example, a compound of which
10 nitrogen atom is oxidized in case when T, U, V or/and W in the formula (I) is or are nitrogen.

"An agent for treatment" refers to a medicament which is employed for the treatment and/or prophylaxis of various diseases.

15 In order to disclose the aforesaid compounds of the general formula (I) of the present invention more specifically, the various symbols used in the formula (I) are explained in more detail by presenting preferred embodiments.

A, B, C and D are independently methine or nitrogen, said
20 methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$, and at least one of A,
25 B, C and D is said methine group.

"Methine which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower

alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$ " refers to unsubstituted methine or methine having a substituent, wherein said substituent may be selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, 5 halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$.

Halogen as said substituent preferably includes fluorine, chlorine and the like.

Lower alkyl as said substituent preferably includes 10 methyl, ethyl and the like.

Halo-lower alkyl as said substituent preferably includes difluoromethyl, trifluoromethyl and the like.

Lower alkoxy as said substituent preferably includes methoxy, ethoxy and the like.

15 Halo-lower alkoxy as said substituent preferably includes difluoromethoxy, trifluoromethoxy and the like.

Lower alkoxycarbonyl as said substituent preferably includes methoxycarbonyl, ethoxycarbonyl and the like.

Lower alkylsulfonyl as said substituent preferably 20 includes methylsulfonyl, ethylsulfonyl and the like.

Lower alkylsulfonyloxy as said substituent preferably includes methylsulfonyloxy, ethylsulfonyloxy and the like.

In a group of formula: $-N(R^1)R^2$ as said substituent, R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , 25 taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino.

Lower alkyl as R^1 or R^2 preferably includes methyl, ethyl, propyl and the like.

"Lower alkylene which may be intervened by oxygen, sulfur

or imino" formed by taking R^1 and R^2 together preferably includes pentamethylene, 3-oxapentamethylene and the like, and it, together with nitrogen next thereto, forms piperidino, morpholino, etc.

5 The preferred embodiment of R^1 and R^2 includes the case where at least one of R^1 or R^2 is lower alkyl, or the case where R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino.

 Thus, a group of formula: $-N(R^1)R^2$ includes, for example,
10 amino, methylamino, ethylamino, propylamino, dimethylamino, diethylamino, ethylmethylamino, 1-azetidiny, 1-pyrrolidinyl, piperidino, morpholino, thiomorpholino, 1-piperazinyl and the like, among which the preferred are methylamino, dimethylamino, piperidino, morpholino and the like.

15 In a group of formula: $-Q^1-Ar^1$ as said substituent, Ar^1 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy,
20 halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxy-carbonyl, lower alkanoylamino and $-Q^2-Ar^2$; Q^1 is a single bond, oxygen, carbonyl or $-N(R^3)-$.

 "Aryl or heteroaryl, any of which is optionally
25 substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxy-carbonyl, lower

alkanoylamino and $-Q^2-Ar^2$ refers to unsubstituted said aryl or said heteroaryl, or said aryl or said heteroaryl having substituent(s) at the substitutable, arbitrary position(s) wherein said substituent(s) may be one, two or more, preferably
5 one or two member(s) identically or differently selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower
10 alkoxycarbonyl, lower alkanoylamino and $-Q^2-Ar^2$.

Halogen as said substituent preferably includes fluorine, chlorine, bromine and the like.

Lower alkyl as said substituent preferably includes methyl, ethyl, propyl, isopropyl and the like.

15 Halo-lower alkyl as said substituent preferably includes difluoromethyl, trifluoromethyl and the like.

Hydroxy-lower alkyl as said substituent preferably includes hydroxymethyl, 2-hydroxyethyl, 1-hydroxy-1-methylethyl and the like.

20 Cyclo-lower alkyl as said substituent preferably includes cyclopropyl, cyclobutyl and the like.

Lower alkenyl as said substituent preferably includes vinyl, 1-propenyl, 2-methyl-1-propenyl and the like.

Lower alkoxy as said substituent preferably includes
25 methoxy, ethoxy and the like.

Halo-lower alkoxy as said substituent preferably includes fluoromethoxy, difluoromethoxy, trifluoromethoxy and the like.

Lower alkylthio as said substituent preferably includes

methylthio, ethylthio and the like.

Lower alkylsulfonyl as said substituent preferably includes methylsulfonyl, ethylsulfonyl, propylsulfonyl and the like.

5 Lower alkanoyl as said substituent preferably includes acetyl, propionyl and the like.

Lower alkoxycarbonyl as said substituent preferably includes methoxycarbonyl, ethoxycarbonyl and the like.

10 Lower alkanoylamino as said substituent preferably includes acetylamino, propanoylamino and the like.

In a group of formula: $-Q^2-Ar^2$ as said substituent, Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, 15 hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl; Q^2 is a single bond, oxygen, carbonyl or $-N(R^3)-$.

"Aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl" refers to unsubstituted said aryl or said heteroaryl, or said aryl or said heteroaryl having substituent(s) at the 20 substitutable, arbitrary position(s) wherein said substituent(s) may be one, two or more, preferably one or two member(s) identically or differently selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, 25

lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl.

Halogen as said substituent preferably includes fluorine, chlorine and the like.

Lower alkyl as said substituent preferably includes
5 methyl, ethyl, propyl, isopropyl and the like.

Halo-lower alkyl as said substituent preferably includes difluoromethyl, trifluoromethyl and the like.

Hydroxy-lower alkyl as said substituent preferably includes hydroxymethyl, 2-hydroxyethyl,
10 1-hydroxy-1-methylethyl and the like.

Lower alkoxy as said substituent preferably includes methoxy, ethoxy and the like.

Halo-lower alkoxy as said substituent preferably includes fluoromethoxy, difluoromethoxy, trifluoromethoxy and
15 the like.

Lower alkylamino as said substituent preferably includes methylamino, ethylamino and the like.

Di-lower alkylamino as said substituent preferably includes dimethylamino, diethylamino and the like.

20 Lower alkanoyl as said substituent preferably includes acetyl, propionyl and the like.

Aryl as said substituent preferably includes phenyl and the like.

The substituent(s) of Ar² preferably include(s) halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl,
25 hydroxy, halo-lower alkoxy and the like.

Aryl as Ar² preferably includes phenyl and the like, and heteroaryl as Ar² preferably includes pyridyl, quinolyl and the like.

In the group of formula: $-N(R^3)-$ as Q^1 and Q^2 , R^3 is hydrogen or lower alkyl.

R^3 preferably includes hydrogen, methyl, ethyl and the like.

5 Q^2 preferably includes a single bond and the like.

The substituent(s) of Ar^1 include(s) preferably halogen, hydroxy, lower alkyl, halo-lower alkyl, lower alkenyl, lower alkoxy, lower alkanoyl, $-Q^2-Ar^2$ and the like, more preferably halogen, hydroxy, halo-lower alkyl, lower alkoxy and the like.

10 Aryl as Ar^1 preferably includes phenyl and the like, and heteroaryl as Ar^1 includes preferably imidazolyl, furyl, thienyl, pyrazolyl, thiazolyl, oxazolyl, isoxazolyl, 1,2,4-oxadiazol, 1,3,4-oxadiazol, pyridyl, pyrazinyl, pyrimidinyl, benzofuranyl, quinolyl and the like, more
15 preferably pyridyl, quinolyl and the like, most preferably pyridyl and the like.

Consequently, Ar^1 includes, for example, phenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,3-difluorophenyl, 2,4-difluorophenyl, 2,6-difluorophenyl,
20 3,5-difluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2-chloro-4-fluorophenyl, 2-chloro-5-fluorophenyl, 2-chloro-6-fluorophenyl, 2-bromo-4-fluorophenyl, 2-bromo-5-fluorophenyl, 2-methylphenyl, 3-methylphenyl,
25 4-methylphenyl, 5-fluoro-2-methylphenyl, 3-fluoromethylphenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 5-fluoro-2-methoxyphenyl, 3-fluoromethoxyphenyl,

2-difluoromethoxyphenyl, 3-difluoromethoxyphenyl,
 2-hydroxy-4-fluorophenyl, 2-hydroxymethylphenyl,
 3-hydroxymethylphenyl, 3-hydroxyphenyl, 4-hydroxyphenyl,
 2-imidazolyl, 2-furyl, 2-thienyl, 1,2,4-oxadiazol-5-yl,
 5 1,3,4-oxadiazol-2-yl, 1,2,4-thiadiazol-5-yl,
 1,3,4-thiadiazol-2-yl, 2-pyridyl, 3-pyridyl, 4-pyridyl,
 2-fluoro-5-pyridyl, 3-fluoro-6-pyridyl, 2-pyrimidinyl,
 4-benzo[b]furanyl, 5-benzo[b]furanyl, 7-benzo[b]furanyl,
 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl,
 10 8-quinolyl and the like. Preferably, Ar¹ includes phenyl,
 2-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl,
 2,5-difluorophenyl, 2-chloro-4-fluorophenyl,
 2-chloro-5-fluorophenyl, 2-trifluoromethylphenyl,
 2-difluoromethoxyphenyl, 3-difluoromethoxyphenyl,
 15 3-hydroxyphenyl, 4-hydroxyphenyl, 2-pyridyl,
 2-fluoro-5-pyridyl, 3-fluoro-6-pyridyl, 7-benzo[b]furanyl,
 2-quinolyl, 3-quinolyl and the like.

Q¹ includes preferably a single bond, oxygen, -N(R³)- and the like, more preferably a single bond and the like.

20 The substituent(s) of methine as A, B, C or D include(s) preferably halogen, halo-lower alkyl, lower alkoxy carbonyl, -N(R¹)R², -Q¹-Ar¹ and the like, more preferably -Q¹-Ar¹ and the like.

The preferred embodiment of A, B, C and D includes, for
 25 example, the case where A and D are same or differently unsubstituted methine or nitrogen, and one of B and C is methine having -Q¹-Ar¹ and the other is unsubstituted methine or nitrogen, more preferably the case where A is unsubstituted methine, B and/or D are nitrogen, and C is methine having -Q¹-Ar¹.

E is methine, hydroxy substituted methine or nitrogen, preferably methine.

n is 0 or 1, preferably 0.

T, U, V and W are independently methine or nitrogen, said
5 methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group.

"Methine which is optionally substituted by a
10 substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy" refers to unsubstituted methine or methine having a substituent, wherein said substituent may be selected from the group consisting of halogen, lower alkyl, halo-lower
15 alkyl, hydroxy, lower alkoxy and halo-lower alkoxy.

Halogen as said substituent preferably includes fluorine, chlorine and the like.

Lower alkyl as said substituent preferably includes methyl, ethyl and the like.

20 Halo-lower alkyl as said substituent preferably includes difluoromethyl, trifluoromethyl and the like.

Lower alkoxy as said substituent preferably includes methoxy, ethoxy and the like.

Halo-lower alkoxy as said substituent preferably
25 includes fluoromethoxy, difluoromethoxy, trifluoromethoxy and the like.

Said substituent includes preferably halogen, lower alkyl, hydroxy, lower alkoxy and the like, more preferably halogen and the like.

The preferred embodiment of T, U, V and W includes, for example, the case where T, U, V and W are independently methine optionally having said substituent, preferably halogen, lower alkyl, hydroxy and lower alkoxy, more preferably halogen; or
5 the case where one of T, U, V and W is nitrogen, preferably the case where one of T, U, V and W is nitrogen and one of the remainder is methine having said substituent, preferably halogen, lower alkyl, hydroxy and lower alkoxy, more preferably halogen; or the case where one of T, U, V and W is nitrogen and
10 all of the remainder are unsubstituted methine.

X is $-\text{N}(\text{SO}_2\text{R}^4)-$, $-\text{N}(\text{COR}^5)-$ or $-\text{CO}-$. Y is $-\text{C}(\text{R}^6)(\text{R}^7)-$, $-\text{O}-$ or $-\text{N}(\text{R}^8)-$. R^4 is lower alkyl, aralkyl or aryl. R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl. R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl
15 or aryl.

Lower alkyl as R^4 , R^5 , R^6 , R^7 or R^8 , each independently, preferably includes methyl, ethyl, propyl and the like.

Aralkyl as R^4 , R^5 , R^6 , R^7 or R^8 , each independently, preferably includes benzyl and the like.

20 Aryl as R^4 , R^5 , R^6 , R^7 or R^8 , each independently, preferably includes phenyl and the like.

R^4 and R^5 preferably include, for example, lower alkyl and the like.

The preferred embodiment of R^6 and R^7 includes, for example,
25 the case where both R^6 and R^7 are hydrogen.


R^8 includes preferably, for example, hydrogen, lower alkyl and the like, more preferably hydrogen and the like.

The preferred embodiment of n, X and Y includes, for example, the case where n is 0, X is $-\text{N}(\text{SO}_2\text{R}^4)-$ or $-\text{N}(\text{COR}^5)-$,

preferably $-\text{N}(\text{SO}_2\text{R}^4)-$, and Y is $-\text{C}(\text{R}^6)(\text{R}^7)-$; or the case where n is 0 or 1, preferably 0, X is $-\text{CO}-$, and Y is $-\text{O}-$ or $-\text{N}(\text{R}^8)-$, preferably $-\text{O}-$. In the latter case, it is more preferable when X is $-\text{CO}-$ and Y is $-\text{O}-$ or $-\text{NH}-$, and further preferable when X is $-\text{CO}-$ and Y is $-\text{O}-$.







The present invention excludes a compound of the formula (I), wherein at the same time E is nitrogen, n is 0, X is $-\text{CO}-$ and Y is $-\text{O}-$.

In more detail, a group of formula (a):




(a)

in the formula (I) includes a group of formula (a₁):

(a₁)


wherein methine in the formula (a₁) is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower

alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$, and the like, more preferably a group of formula (a₂)




(wherein Ar^1 and Q^1 have the aforesaid meaning), and the like.

5 Moreover, a group of formula (b):



in the formula (I) includes a group of formula (b₁):


27



(wherein R⁰ is hydrogen or hydroxy; R⁴ has the aforesaid meaning),


5 and the like.

Preferred compounds of the formula (I) are, for example, compounds of the formula (I-a):




(wherein A, B, C, D, E and R⁴ have the aforesaid meaning);

5 compounds of the formula (I-b):



(wherein A, B, C, D, R⁰, T, U, V and W have the aforesaid meaning);

compounds of the formula (I-c):



10 (wherein A, B, C, D and E have the aforesaid meaning);
or the like.


Among the compounds of the formula (I), the formula (I-a), the formula (I-b) or the formula (I-c), the preferred compounds are, for example, compounds in which A, B, C and D are aforesaid
15 preferred embodiment, Ar¹ is phenyl or heteroaryl, more preferably phenyl, any of which is optionally substituted by a substituent selected from the group consisting of halogen,


nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsufonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino and $-Q^2-Ar^2$, and R^0 is hydrogen.

Further, among the compound of the formula (I-b), the preferred compounds are, for example, compounds in which all of T, U, V and W are unsubstituted methine; or compounds in which one of T, U, V and W is nitrogen, preferably one of T, U, V and W is nitrogen and all of the remainder are unsubstituted methine or one of the remainder is methine having fluorine or chlorine.



The compounds of the present invention may include stereoisomers such as optical isomers, diastereoisomers and geometrical isomers, or tautomers depending upon the mode of substituents. The compounds of the present invention include all the stereoisomers, tautomers and their mixtures.

For example, compounds of the formula (I-a) include stereoisomers such as a trans-form compound of the formula (I-1a) or (I-1a'):






and a cis-form compound of the formula (I-2a) or (I-2a'):




5

, and compounds of the formulae (I-1a) and (I-2a') are preferable.



Compounds of the formula (I-b) include stereoisomers such as a trans-form compound of the formula (I-1b) or (I-1b'):



10





and a cis-form compound of the formula (I-2b) or (I-2b'):





5 , and compounds of formulae (I-1b) and (I-2b') are preferable.

Compounds of the formula (I-c) wherein E is methine, include stereoisomers such as a trans-form compound of the formula (I-1c) or (I-1c'):





and a cis-form compound of the formula (I-2c) or (I-2c'):



5 , and compounds of the formulae (I-1c) and (I-2c') are preferable.

Also included within the scope of the invention are polymorphs, hydrates and solvates of the compounds of the present invention.

10 The present invention also includes prodrugs of the compounds of this invention within its scope. In general, such prodrugs are functional derivatives of the compounds of the present invention which can be readily converted *in vivo* into the required compound. Thus, in the treatment methods for
 15 various diseases according to the present invention, the term "administering" shall encompass not only administration of the compound specified in this disclosure but also administration

of a compound which is converted *in vivo* into the specified compound when it is administered to a patient. Conventional procedures for selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs," ed. H. Bundgaard, Elsevier (1985), which are
5 entirely incorporated by reference in this specification. Metabolites of these compounds include active compounds produced upon introduction of compounds of the present invention into the biological milieu, and are encompassed in
10 the scope of the present invention.

The specific compound of the formula (I) is, for example,
2-[1-methylsulfonylspiro[indoline-3,4'-piperidin]-1'-yl]-5-chlorobenzimidazole,
5,6-dichloro-2-[1-methylsulfonylspiro[indoline-3,4'-
15 piperidin]-1'-yl]benzimidazole,
5-chloro-2-[1-ethylsulfonylspiro[indoline-3,4'-piperidin]-1'-yl]benzimidazole,
trans-5-chloro-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-6-(trifluoromethyl)benzimidazole,
20 trans-5-(4-fluorophenyl)-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]benzimidazole,
trans-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-5-phenylbenzimidazole,
trans-5-(3-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-
25 1(3H),1'-cyclohexan]-4'-yl]benzimidazole,
trans-5-(4-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]benzimidazole,
trans-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-5-(2-pyridyl)imidazo[4,5-b]pyridine,

- trans-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-6-(2-pyridyl)imidazo[4,5-b]pyridine,
trans-5-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-b]pyridine,
5 trans-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-6-phenylimidazo[4,5-c]pyridine,
trans-6-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-c]pyridine,
trans-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-5-phenylimidazo[4,5-b]pyrazine,
10 trans-5-(4-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-b]pyrazine,
trans-6-(4-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-c]pyridazine,
15 trans-2-(2-fluorophenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(3-fluorophenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(4-fluorophenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
20 trans-2-(2-methoxyphenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(4-methoxyphenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
25 trans-2-(2-methylphenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-(2-trifluoromethylphenyl)purine,
trans-2-(2-chloro-4-fluorophenyl)-8-[3'-oxospiro[cyclo-

- hexane-1,1'-(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-hydroxymethylphenyl)-8-[3'-oxospiro[cyclo-
hexane-1,1'-(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-furyl)-8-[3'-oxospiro[cyclohexane-1,1'-(3'H)-
5 isobenzofuran]-4-yl]purine,
trans-8-[3'-oxospiro[cyclohexane-1,1'-(3'H)-isobenzofuran]-
-4-yl]-2-(2-thienyl)purine,
trans-8-[3'-oxospiro[cyclohexane-1,1'-(3'H)-isobenzofuran]-
-4-yl]-2-(2-pyrrolyl)purine,
10 trans-2-(3-fluoropyridin-6-yl)-8-[3'-oxospiro[cyclohexane-
1,1'-(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-chloropyridin-6-yl)-8-[3'-oxospiro[cyclohexane-
1,1'-(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-fluorophenoxy)-8-[3'-oxospiro[cyclohexane-
15 1,1'-(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2,6-difluorophenoxy)-8-[3'-oxospiro[cyclohexane-
1,1'-(3'H)-isobenzofuran]-4-yl]purine,
trans-8-[3'-oxospiro[cyclohexane-1,1'-(3'H)-isobenzofuran]-
4-yl]-2-(1-piperidyl)purine,
20 trans-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclo-
hexan]-4'-yl]-2-phenylpurine,
trans-2-(2-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(3-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
25 1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2-chlorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine,

- trans-2-(3-chlorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- trans-2-(4-chlorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- 5 trans-2-(2-chloro-4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- trans-2-(4-methoxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- trans-2-(2-methylphenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- 10 trans-2-(2-difluoromethoxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- trans-2-(3-difluoromethoxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- 15 trans-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-2-(2-trifluoromethylphenyl)purine,
- trans-2-(2,4-difluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- trans-2-(2,5-difluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- 20 trans-2-(2-bromo-4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- trans-2-(4-chloro-2-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
- 25 trans-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-2-(3-quinolyl)purine,
- trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-2-phenylpurine,
- trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-

- cyclohexan]-4'-yl]-2-(2-fluorophenyl)purine,
trans-2-(2,4-difluorophenyl)-8-[5-fluoro-3-oxospiro[6-aza-
isobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2,5-difluorophenyl)-8-[5-fluoro-3-oxospiro[6-aza-
5 isobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-8-[7-hydroxy-3-oxospiro[6-aza-
isobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
5,6-dichloro-2-[3,4-dihydro-3-oxospiro[isoquinoline-
1(2H),4'-piperidin]-1'-yl]benzimidazole,
10 2-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-
1'-yl]-5-phenylbenzimidazole,
8-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-
1'-yl]-2-phenylpurine,
8-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-
15 1'-yl]-2-(2-fluorophenyl)purine,
8-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-
1'-yl]-2-(4-fluorophenyl)purine,
trans-8-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-
isoquinolin]-4-yl]-2-phenylpurine,
20 trans-2-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-
isoquinolin]-4-yl]-5-phenylbenzimidazole,
trans-8-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-
isoquinolin]-4-yl]-2-(2-fluorophenyl)purine,
trans-8-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-
25 isoquinolin]-4-yl]-2-(4-fluorophenyl)purine,
trans-2-(4-fluorophenyl)-8-[7-hydroxy-3-oxospiro[6-azaiso-
benzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine 6-oxide,

trans-2-(4-fluoro-2-hydroxyphenyl)-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-6-hydroxy-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine,
5 trans-2-(4-hydroxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H), 1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluoro-3-hydroxyphenyl)-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine,
cis-2-(4-fluorophenyl)-8-[4'-hydroxy-3-oxospiro[6-azaiso-
10 benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine,
and the like.

Among these compounds, the preferable compound is, for example,

trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-
15 4-yl]-2-phenylpurine,
trans-2-(2-fluorophenyl)-8-[3'-oxospiro[cyclohexane-
1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H), 1'-cyclohexan]-4'-yl]purine,
20 trans-2-(2,5-difluorophenyl)-8-[3-oxospiro[6-azaisobenzo-
furan-1(3H), 1'-cyclohexan]-4'-yl]purine,
trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H), 1'-
cyclohexan]-4'-yl]-2-(2-fluorophenyl)purine,
trans-5-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-
25 1(3H), 1'-cyclohexan]-4'-yl]imidazo[4, 5-b]pyridine,
and the like.

The process for producing compounds of the present invention is illustrated as follows.


Compounds (I) of the present invention can be synthesized,

for example, by the following production methods or the processes shown in examples, but these embodiments are not intended to restrict the process for producing compounds (I) of this invention.


5

Production Process 1

A compound of the formula (I-1) or N-oxide derivative thereof:



10 (wherein A, B, C, D, n, T, U, V, W, X and Y have each the same meaning as defined above) can be prepared by reacting a compound of the formula (II):



(wherein a, b, c and d are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R^{1p})R^{2p}, -Q^{1p}-Ar^{1p} and optionally protected hydroxy, and at least one of a, b, c and d is said methine;

15

20

Ar^{1p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting

of halogen, nitro, lower alkyl, halo-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, lower alkanoyl, lower alkoxy carbonyl, lower alkanoylamino, $-Q^{2p}-Ar^{2p}$, optionally
 5 protected hydroxy, optionally protected hydroxy-lower alkyl, and optionally protected carboxyl;

Ar^{2p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy,
 10 halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl, aryl, optionally protected hydroxy-lower alkyl and optionally protected hydroxy;

L^1 is a leaving group;


P is an imino-protecting group;

15 Q^{1p} and Q^{2p} are independently a single bond, oxygen, optionally protected carbonyl or $-N(R^3)-$;

R^{1p} and R^{2p} are independently an amino-protecting group, an imino-protecting group, hydrogen or lower alkyl, or R^{1p} and R^{2p} , taken together, form lower alkylene which may be intervened
 20 by oxygen, sulfur or optionally protected imino;

R^3 has the same meaning as defined above)


with a compound of the formula (III):



(III)

(wherein n, X and Y have each the same meaning as defined above,
 25 provided that the compound (III) when n is 0, X is $-CO-$ and Y

is -O- is excluded) or its salt to give a compound of the formula (IV-1):



(wherein a, b, c, d, n, P, t, u, v, w, X and Y have each the same meaning as defined above), and optionally removing wherefrom protecting group(s) and/or oxidizing nitrogen atom(s) of the compound (IV-1).

The leaving groups represented by L^1 include, for example, halogen (e.g. chlorine, bromine, iodine), organic sulfonyl (e.g. methanesulfonyl, ethanesulfonyl, benzenesulfonyl), organic sulfonyloxy (e.g. methanesulfonyloxy, trifluoromethanesulfonyloxy, p-toluenesulfonyloxy) and the like.

The present process refers to a process for preparing a compound of the formula (I) wherein E is nitrogen, namely a compound of the formula (I-1).

In the above reaction, when a reactant has an amino, imino, hydroxy, carboxyl, carbonyl or the like which does not participate in the reaction, the reaction may be carried out after protecting the amino, imino, hydroxy, carboxyl, carbonyl with an amino- or imino-protecting group, a hydroxy-protecting group, a carboxyl-protecting group, or a carbonyl-protecting group, followed by deprotection after completion of the reaction.

The "amino- or imino-protecting group" is not particularly restricted, so long as it has such protective function. There are employed, for example, aralkyl (e.g. benzyl, p-methoxybenzyl, 3,4-dimethoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, benzhydryl, trityl); lower alkanoyl (e.g. formyl, acetyl, propionyl, butyryl, pivaloyl); benzoyl; arylalkanoyl (e.g. phenylacetyl, phenoxyacetyl); lower alkoxycarbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propyloxycarbonyl, tert-butoxycarbonyl); aralkyloxycarbonyl (e.g. benzyloxycarbonyl, p-nitrobenzyloxycarbonyl, phenethyloxycarbonyl); lower alkylsilyl (e.g. trimethylsilyl, tert-butyldimethylsilyl); tetrahydropyranyl; trimethylsilylethoxymethyl; lower alkylsulfonyl (e.g. methylsulfonyl, ethylsulfonyl); arylsulfonyl (e.g. benzenesulfonyl, toluenesulfonyl) and the like, among which the particularly preferred are acetyl, benzoyl, tert-butoxycarbonyl, trimethylsilylethoxymethyl, methylsulfonyl and the like.

The "hydroxy-protecting group" is not particularly restricted, so long as it has such protective function for hydroxy groups. There are employed, for example, lower alkyl (e.g. methyl, ethyl, propyl, isopropyl, tert-butyl); lower alkylsilyl (e.g. trimethylsilyl, tert-butyldimethylsilyl); lower alkoxymethyl (e.g. methoxymethyl, 2-methoxyethoxymethyl); tetrahydropyranyl; trimethylsilylethoxymethyl; aralkyl (e.g. benzyl, p-methoxybenzyl, 2,3-dimethoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, trityl); and acyl (e.g. formyl, acetyl), among which the particularly preferred are methyl, methoxymethyl,

tetrahydropyranyl, trityl, trimethylsilylethoxymethyl, tert-butyldimethylsilyl, acetyl and the like.

The "carboxyl- protecting group" is not particularly restricted, so long as it has such protective function for carboxyl groups. There are employed, for example, lower alkyl (e.g. methyl, ethyl, propyl, isopropyl, tert-butyl); halo-lower alkyl (e.g. 2,2,2-trichloroethyl); lower alkenyl (e.g. 2-propenyl); aralkyl (e.g. benzyl, p-methoxybenzyl, p-nitrobenzyl, benzhydryl, trityl) and the like, among which the particularly preferred are methyl, ethyl, tert-butyl, 2-propenyl, benzyl, p-methoxybenzyl or benzhydryl and the like.

The "carbonyl-protecting group" is not particularly limited, so long as it has such protective function for carbonyl groups. There are employed, for example, acetals or ketals, such as ethylene ketal, trimethylene ketal, and dimethyl ketal and the like.

The reaction between a compound of the formula (II) and a compound of the formula (III) is usually carried out by employing an equivalent to excessive mole, preferably an equivalent to 1.5 moles, of compound (III), relative to 1 mole of compound (II).

The reaction is usually carried out in an inert solvent. Preferable examples of such solvent are methylene chloride, chloroform, tetrahydrofuran, dioxane, dimethylformamide, N-methylpyrrolidone, dimethyl sulfoxide and the like, or a mixture thereof and the like.

The reaction is preferably carried out in the presence of a base such as organic bases (e.g. triethylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine),

inorganic bases (e.g. sodium hydrogen carbonate, sodium carbonate, potassium carbonate, cesium carbonate, sodium hydroxide, potassium hydroxide) and the like.

The base is used in equimolar amount or in excess mole,
5 preferably 1 to 5 moles, relative to 1 mole of the compound of the formula (II).

The reaction temperature is usually from 0°C to 200°C, preferably 20°C to 150°C.

The reaction time is usually 5 minutes to 7 days,
10 preferably 30 minutes to 24 hours.

A usual treatment is carried out after completion of the reaction to obtain a crude product of a compound of the formula (IV-1). The resulting compound of the formula (IV-1) is, with or without purification according to the common method,
15 subjected to optional removal of the protecting group for the amino, hydroxy, carboxyl, carbonyl or the like, thereby a compound of the formula (I-1) can be prepared.

Although the method for the removal of said protecting groups depends upon the kinds of the protecting groups, the
20 stability of a desired compound (I-1) and the like, it is carried out by, for example, a solvolysis using an acid or a base, that is, a method wherein for example 0.01 mole to a large excess of acid, preferably trifluoroacetic acid, formic acid, hydrochloric acid and the like, or an equivalent mole to a large
25 excess of base, preferably potassium hydroxide, calcium hydroxide and the like is acted; a chemical reduction using a metal hydride complex; or a catalytic reduction using a palladium-carbon catalyst, a Raney-nickel catalyst, etc.; and the like, according to, for example, a method described in the

literature (Protective Groups in Organic Synthesis, T.W.Greene, John Wiley & Sons, (1981)) or its similar method.

The oxidation of a nitrogen atom may be carried out by using of an oxydizing agent (for example m-chloroperbenzoic acid, dioxirane, sodium periodate and hydrogen peroxide).
 5 Reaction between a compound of the general formula (IV-1) and an oxydizing agent is usually carried out by employing 0.5 mole to excessive moles, preferably 1 mole to 5 moles of the oxydizing agent based on 1 mole of compound (IV-1).


10 The reaction is usually carried out in an appropriate solvent which depend on the oxydizing agent used in the reaction. Preferable examples of the solvent include methylene chloride and chloroform for m-chloroperbenzoic acid, acetone and water for dioxirane.

15 Reaction temperature is usually -50 °C to 100 °C, preferably -20 °C to 50 °C.

Reaction time is usually 15 minutes to 7 days, preferably 30 minutes to 24 hours.


20 Production Process 2

A compound of the formula (I-2) or N-oxide derivative thereof:




(wherein A, B, C, D, n, R⁰, T, U, V, W, X and Y have each the
 25 same meaning as defined above) can be prepared by reacting a

compound of the formula (V):




(wherein a, b, c and d have each the same meaning as defined above) or its salt with a compound of the formula (VI):




5

(wherein R^{Op} is hydrogen or optionally protected hydroxy; n, t, u, v, w, X and Y have each the same meaning as defined above) or its salt to give a compound of the formula (VII):



10 (wherein a, b, c, d, n, R^{Op} , t, u, v, w, X and Y have each the same meaning as defined above), followed by subjecting the compound (VII) to intramolecular dehydrative ring closure to give a compound of the formula (VIII):



15 (wherein a, b, c, d, n, R^{Op} , t, u, v, w, X and Y have each the

same meaning as defined above), and optionally removing wherefrom the protecting group(s) and/or oxidizing nitrogen atom(s) of the compound (VIII).

The present process refers to a process for preparing a
5 compound of the formula (I) wherein E is methine or hydroxy substituted methine, namely a compound of the formula (I-2).

The reaction between a compound of the formula (V) and a carboxylic acid of the formula (VI) is usually carried out by employing 0.5 moles to excessive mole, preferably 1 mole to
10 1.5 moles of the carboxylic acid (VI), relative to 1 mole of a compound of the formula (V).

The reaction is usually carried out in an inert solvent. Preferable examples of such solvent are methylene chloride, chloroform, tetrahydrofuran, dimethylformamide, pyridine and
15 the like, and a mixture thereof and the like.

The above reaction is preferably carried out in the presence of a condensing agent including, for example, N,N'-dicyclohexylcarbodiimide, N,N'-diisopropylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide,
20 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, benzotriazol-1-yloxy-tris-(dimethylamino)phosphonium hexafluorophosphate, benzotriazol-1-yloxy-tris-pyrrolidino-phosphonium hexafluorophosphate, bromotris(dimethylamino)phosphonium
25 hexafluorophosphate, diphenylphosphoric azide, 1,1'-carbonyldiimidazole and the like.

Such condensing agent can be usually used in 1 mole to excess moles, preferably 1 to 1.5 moles, relative to 1 mole of a compound of the formula (VI).

The reaction temperature is usually from -50°C to 100°C, preferably -20°C to 50°C.

The reaction time is usually 30 minutes to 7 days, preferably 1 to 24 hours.

5 In place of the carboxylic acid of the formula (VI), an activated derivative of said carboxylic acid may be reacted with a compound of the formula (V), thereby a compound of the formula (I-2) can be prepared.

10 Examples of such activated derivatives of the carboxylic acid of the formula (VI) are acid halides, mixed anhydrides, active esters, active amides and the like.

15 The acid halides of the carboxylic acid of the formula (VI) can be prepared by reacting a carboxylic acid of the formula (VI) with a halogenating agent in a conventional manner. The halogenating agent used includes, for example, thionyl chloride, phosphorus trichloride, phosphorus pentachloride, phosphorus oxychloride, phosphorus tribromide, oxalyl chloride, phosgene and the like.

20 The mixed anhydrides of the carboxylic acid of the formula (VI) can be prepared by reacting a carboxylic acid of the formula (VI) with an alkyl chlorocarbonate (e.g. ethyl chlorocarbonate), an aliphatic carboxylic acid chloride (e.g. pivaloyl chloride) and the like according to the conventional method.

25 The active esters of the carboxylic acid of the formula (VI) can be prepared by reacting a carboxylic acid of the formula (VI) with an N-hydroxy compound (e.g. N-hydroxysuccinimide, N-hydroxyphthalimide, 1-hydroxybenzotriazole); a phenol compound (e.g. 4-nitrophenol, 2,4-dinitrophenol, 2,4,5-trichlorophenol, pentachlorophenol) and the like in the

presence of a condensing agent (e.g. N,N'-dicyclohexyl-carbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide) according to the conventional method.

The active amides of the carboxylic acid of the formula
5 (VI) can be prepared by reacting a carboxylic acid of the formula (VI) with 1,1'-carbonyldiimidazole, 1,1'-carbonylbis(2-methyl-imidazole) and the like according to the conventional method.

The reaction between a compound of the formula (V) and
10 a reactive derivative of the carboxylic acid of the formula (VI) is usually carried out by employing 0.5 moles to excessive mole, preferably 1 mole to 1.5 moles of the reactive derivative of the carboxylic acid (VI), relative to 1 mole of compound (V).

The reaction is usually carried out in an inert solvent.
15 Preferable examples of such inert solvent are methylene chloride, chloroform, tetrahydrofuran, dimethylformamide, pyridine and the like, and a mixture thereof and the like.

The above reaction may proceed in the absence of a base, but it is preferable to carry out the reaction in the presence
20 of a base to promote the reaction smoothly.

The bases include organic bases (e.g. triethylamine, diisopropylethylamine, pyridine, 4-dimethylaminopyridine), or inorganic bases (e.g. sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydrogen
25 carbonate).

It is preferable to employ 1 mole to excessive mole of said base to 1 mole of a compound of the formula (V). When the base is liquid, such base can also be used as a solvent.

The reaction temperature is usually -50°C to 100°C.

preferably -20°C to 50°C.

The reaction time is usually 5 minutes to 7 days, preferably 30 minutes to 24 hours.

A usual treatment is carried out after completion of the
5 reaction to obtain a crude product of a compound of formula (VII). The resulting compound of the formula (VII) may be, with or without purification according to the conventional manner, subjected to optional intramolecular ring closure condensation.

10 The intramolecular ring closure condensation for preparing a compound of the formula (VIII) from the compound (VII) is usually carried out in the presence of an inert solvent or without any solvent.

Preferred examples of such inert solvents are ethanol,
15 propanol, butanol, pentanol, 1,4-dioxane, dimethoxyethane, dimethylformamide, dimethyl sulfoxide, benzene, toluene, xylene, pyridine and the like, and a mixture thereof and the like.

The reaction temperature is usually from room temperature
20 to the boiling point of the solvent used, preferably 80°C to 190°C.

The reaction time is usually from 5 hours to 7 days, preferably from 12 hours to 3 days.

The above ring closure may be carried out in the presence
25 of a dehydrating agent or a catalytic-amount of Lewis acid. The dehydrating agent includes, for example, cesium fluoride, phosphorus oxychloride, phosphorus pentachloride, polyphosphoric acid, thionyl chloride and the like. As the Lewis acid, there are exemplified by scandium

trifluoromethanesulfonate, yttrium trifluoromethanesulfonate, lanthanum trifluoromethanesulfonate, lanthanide trifluoromethanesulfonate and the like. The ring closure is carried out preferably without any solvent, or in the presence
5 of a solvent such as methylene chloride, chloroform, benzene, toluene, xylene and the like or a mixture thereof.

The amount of the dehydrating agent to be used is usually 1 mole to excessive mole, preferably 2 to 10 moles, relative to 1 mole of a compound of the formula(VII), and that of the
10 Lewis acid is 10 to 200 mole %, preferably 10 to 100 mole %.

In general, the reaction temperature is preferably from room temperature to the boiling point of the solvent used.


The reaction time is from one hour to 7 days, preferably from 5 hours to 3 days.

15 A compound of the formula (I-2) can be produced by treating a reaction mixture in the usual way after removal of the said protecting group(s) when the product has a protecting group after completion of the reaction, or by treating the mixture directly in the usual way when the protecting group is absent.


20 The removal of the protecting group(s), the oxidation of nitrogen atom(s) and the post-treatment may be conducted according to the method described in the above Production Process 1.

25 Production Process 3


A compound of the formula (I-2) or N-oxide derivative thereof:




(wherein A, B, C, D, n, R⁰, T, U, V, W, X and Y have each the same meaning as defined above) can be prepared by reacting a compound of the formula (V):



(wherein a, b, c and d have each the same meaning as defined above) or its salt with a compound of the formula (IX):



(wherein n, t, R^{0p}, u, v, w, X and Y have each the same meaning as defined above) or its salt to give a compound of the formula (VIII):



(wherein a, b, c, d, n, R^{0p}, t, u, v, w, X and Y have each the same meaning as defined above), and optionally removing
wherefrom protecting group(s) and/or oxidizing nitrogen

atom(s) of the compound (VIII).

The present process refers to a process for preparing a compound of the formula (I) wherein E is methine or hydroxy substituted methine, namely a compound of the formula (I-2).

5 The reaction between a compound of the formula (V) and a compound of the formula (IX) is usually carried out by employing 0.5 to 5 moles, preferably 0.7 to 3 moles of a compound of the formula (IX), relative to 1 mole of a compound of the formula (V).

10 The reaction is usually carried out in the absence of a solvent or in an inert solvent. Preferred examples of the inert solvent are benzene, toluene, xylene, methylene chloride, chloroform, hexane and the like, and a mixture thereof and the like.

15 The reaction temperature is usually from -20°C to the boiling point of the solvent used, preferably 20°C to 200°C.

The reaction time is usually from 30 minutes to 7 days, preferably from 3 hours to 3 days.

20 The above reaction is preferably carried out in the presence of a Lewis acid such as zinc dichloride, titanium tetrachloride, scandium trifluoromethanesulfonate, ytterbium trifluoromethanesulfonate, lanthanum trifluoromethanesulfonate and the like.

25 The amount of the Lewis acid used is 10 to 200 mole %, preferably 20 to 100 mole % per mole of a compound of the formula (V).

When the reaction is carried out in the presence of a Lewis acid, it is preferable to effect the reaction without any solvent, or in the presence of a solvent such as methylene

chloride, chloroform, benzene, toluene, xylene and the like, or a mixture thereof and the like.

In general, the reaction temperature is from 0°C to the boiling point of the solvent used, preferably room temperature
5 to 150°C.


The reaction time is usually from 1 hour to 7 days, preferably 12 hours to 3 days.

A compound of the formula (I-2) can be produced by treating a reaction mixture in the usual way after removal of the said
10 protecting group(s) when the product has a protecting group after completion of the reaction, or by treating the mixture directly in the usual way when the protecting group is absent.

The removal of the protecting group(s), the oxidation of nitrogen atom(s) and the post-treatment may be conducted
15 according to the method described in the above Production Process 1.


Production Process 4

A compound of the formula (I-3) or N-oxide derivative
20 thereof:




(wherein A¹, B¹, C¹ and D¹ are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower
25 alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower

alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Ar^1$, and at least one of A^1 , B^1 , C^1 and D^1 is said methine having a group represented by $-Ar^1$; Ar^1 , E , n , R^1 , R^2 , T , U , V , W , X and Y have each the same meaning as defined above) can be prepared by reacting a compound of the formula (X):




(wherein a^0 , b^0 , c^0 and d^0 are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^{1p})R^{2p}$ and optionally protected hydroxy, and at least one of a^0 , b^0 , c^0 and d^0 is said methine group having a halogen or trifluoromethanesulfonyloxy group; E^p is nitrogen, methine or methine substituted by optionally protected hydroxy; and n , P , R^{1p} , R^{2p} , t , u , v , w , X and Y have each the same meaning as defined above)

with a compound of the formula (XI):



(wherein Met is a common organometallic atom, and Ar^{1p} has the same meaning as defined above) in the presence of a catalyst to give a compound of the formula (XII):



(wherein a^1 , b^1 , c^1 and d^1 are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^{1p})R^{2p}$, $-Ar^{1p}$ and optionally protected hydroxy, and at least one of a^1 , b^1 , c^1 and d^1 is said methine having a group represented by $-Ar^{1p}$; and Ar^{1p} , E^p , n , P , R^{1p} , R^{2p} , t , u , v , w , X and Y have each the same meaning as defined above), and optionally removing wherefrom the protecting group(s) and/or oxidizing nitrogen atom(s) of the compound (XII).

The present process refers to a process for preparing a compound of the formula (I) wherein at least one of A , B , C and D is methine having a group represented by $-Ar^1$, namely a compound of the formula (I-3).

The common organometallic atoms represented by Met are those usually employed in the cross-coupling reaction, and include, for example, lithium, boron, silicon, magnesium, aluminum, zinc, tin and the like, among which boron, zinc, and tin are preferable. As the concrete embodiment of the practical use of such metal atoms, boron is used in the form of boric acid or boric acid ester, etc., zinc is used in the form of zinc chloride, zinc bromide or zinc iodide, etc., and tin is used in the form of tri-lower alkyl tin, etc.

The reaction between a compound of the formula (X) and a compound of the formula (XI) is usually carried out by employing 0.5 to 5 moles, preferably 0.7 to 3 moles of a compound of the formula (XI), relative to 1 mole of a compound of the formula (X).

The catalyst used in the reaction includes a transition metal generally employed in the cross-coupling reaction such as copper, nickel, palladium and the like. More precisely, preferable examples of such catalyst are tetrakis(triphenylphosphine)palladium(0), palladium(II) acetate, bis(triphenylphosphine) palladium(II) chloride, [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride and the like.

The reaction is usually carried out in an inert solvent. Preferred examples of such inert solvents are water, benzene, toluene, xylene, methylene chloride, chloroform, dimethoxyethane, tetrahydrofuran, dioxane, dimethylformamide and the like, or a mixture thereof and the like.

The reaction temperature is usually from room temperature to the boiling point of the solvent used, preferably 20°C to 200°C.

The reaction time is usually 30 minutes to 7 days, preferably 3 hours to 2 days.

The above reaction is preferably carried out in the presence of a base such as inorganic bases (e.g. sodium hydroxide, potassium hydroxide, sodium hydrogen carbonate, sodium carbonate, potassium carbonate, cesium carbonate), organic bases (e.g. triethylamine, diisopropylamine) and the like.

The amount of the base used is usually 0.5 to 5 moles, preferably 0.7 to 3 moles of said base, relative to 1 mole of a compound of the formula (X).

A compound of the formula (I-3) can be prepared by treating
5 a reaction mixture in the usual way after removal of the said protecting group(s) when the product has a protecting group after completion of the reaction, or by treating the mixture directly in the usual way when the protecting group is absent.

The removal of the protecting group(s), the oxidation
10 of nitrogen atom(s) and the post-treatment may be conducted according to the method described in the above Production Process 1.

The compounds of the formula (I-1), (I-2) or (I-3) may be readily isolated and purified by the conventional separation
15 technique, and examples of such technique are solvent extraction, recrystallization, column chromatography, preparative thin layer chromatography and the like.

These compounds may be converted into the pharmaceutically acceptable salts or esters by the conventional
20 method, and on the contrary, the conversion of the salts or esters into free compounds may also be carried out according to the conventional method.

The compounds of the formulae (II), (III), (V), (VI), (IX) and (XI) are commercially available, or can be prepared
25 according to the common methods or analogous methods thereto, or the methods shown in Examples and Reference Examples, optionally employed in combination.

The salts of compounds of formula (III), (V), (VI) and (IX) refer to the common salts, for example, base addition salt

to carboxyl group when the compound has a carboxyl group, or acid addition salt to amino or basic heterocyclyl when the compound has amino or basic heterocyclyl group(s), and the like.

Aforesaid base addition salts include salts with alkali
5 metals (for example sodium, potassium); alkaline earth metals (for example calcium, magnesium); ammonium or organic amines (for example trimethylamine, triethylamine, dicyclohexylamine, ethanolamine, diethanolamine, triethanolamine, procaine, N,N'-dibenzylethylenediamine), and the like.

10 Aforesaid acid addition salts include salts with inorganic acids (for example hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid, perchloric acid), organic acids (for example maleic acid, fumaric acid, tartaric acid, citric acid, ascorbic acid, trifluoroacetic acid), sulfonic acids (for
15 example methanesulfonic acid, isethionic acid, benzenesulfonic acid, p-toluenesulfonic acid), and the like.

The utility of compounds of the present invention as a medicament is proved by the following pharmacological tests.

20 Pharmacological Test 1 (NPY binding inhibition test)

cDNA sequence encoding human NPY Y5 receptor (c.f. International patent publication number WO96/16542) was cloned into expression vectors pcDNA3, pRc/RSV (made by Invitrogen Inc.) and pCI-neo (made by Promega Inc.). The expression
25 vectors thus obtained were transfected to host cells COS-7, CHO and LM(tk-) (American Type Culture Collection) by cationic lipid method (Proceedings of the National Academy of Sciences of the United States of America, 84: 7413(1987)) to give NPY Y5 receptor expression cells.

A membrane sample prepared from the cells which expressed NPY Y5 receptor was incubated together with a test compound and [¹²⁵I]peptideYY (made by NEN) (20,000cpm) in an assay buffer (25 mM Tris buffer, pH7.4, containing 10 mM magnesium chloride, 1 mM phenylmethylsulfonyl fluoride, 0.1% bacitracin and 0.5% bovine serum albumin) at 25° C for 2 hours, then filtered through a glass filter GF/C and washed with 5 mM Tris buffer (pH7.4) containing 0.3% BSA. The radioactivity of the cake on the glass filter was measured. Non-specific binding was measured in the presence of 1 μM peptide YY, and a 50% Inhibitory Concentration (IC₅₀) of the test compound against specific peptideYY binding was determined (Endocrinology, 131: 2090(1992)). The results are shown in Table 1.

Table 1

Compounds	IC ₅₀ (nM)
Example 6	2.2
Example 8	1.8
Example 13	2.9
Example 17	2.7
Example 20	2.3
Example 27	0.72

As shown above, the compounds of this invention potently inhibited peptideYY (NPY homologue) binding to NPY Y5 receptors.

Pharmacological Test 2 (Antagonistic effect on D-Trp³⁴NPY-induced feeding behavior)

A chronic guide cannula (26 gauge, length 11 mm) was inserted stereotaxically into the third cerebral ventricle of male SD rats (7-8 weeks old, 200-300g) anesthetized with ketamine/xylazine (single intraperitoneal administration of

74 and 11 mg/kg) and fixed by dental resin. The top of the guide cannula was located 2.2mm behind bregma, and 8 mm depth from the skull surface on the median line. After about 1-week recovery period, D-Trp³⁴NPY (1µg/0.4µL/head, synthetic
5 cerebrospinal fluid containing 0.05% bovine serum albumin) was injected into the third ventricle. A test compound suspended in 0.5% aqueous methylcellulose was administered orally 2 hours before the administration of D-Trp³⁴NPY, and the food consumption was measured 2 hours after the administration of
10 D-Trp³⁴NPY.

The results revealed that 10 mg/kg of the compound of this invention significantly suppressed the increase in food consumption induced by D-Trp³⁴NPY (NPY homologue) which was administered to the third ventricle.

15

Pharmacological Test 3 (Pharmacokinetics test)

A test compound was orally or intravenously administered to male SD rats (7-10 weeks old, 200-400g) which abstained from food overnight. About 100 µL of blood was collected from the
20 tail vein at predetermined time, using a heparinized capillary. The blood was centrifuged (4°C, 6,000 r.p.m., 10 minutes) to collect the plasma, to which was added 3-fold amount of ethanol containing an internal standard. The mixture was stirred, allowed to stand at -20°C for 20 minutes, and then centrifuged
25 (4°C, 10,000 r.p.m., 10 minutes). The supernatant was analyzed by LC/MS/MS, and the concentration of the test compound in the plasma was measured using a relative calibration curve.

The results revealed that the bioavailability of the compound of Example 6 was 78% and its half-life in the blood

was 3.0 hours.

Pharmacological Test 4 (Brain/cerebrospinal fluid transport test)

5 A test compound was orally or intravenously administered to male SD rats (7-10 weeks old, 200-400g), and whole blood was collected from the abdominal aorta of said rats anesthetized with ether at predetermined time, using a heparin-treated syringe. Then, the head skin was cut open, and a dental 30G
10 needle was inserted between the cervical vertebrae, and it was further inserted into the cavum subarachnoideale. After 50 to 100 μ L cerebrospinal fluid had been collected by a 1 ml-syringe through a tube connected to said dental 30G needle, the brain was extracted. The blood sample was centrifuged (4°C, 6,000
15 r.p.m., 10 minutes) to collect the plasma, to which was added 3-fold amount of ethanol containing an internal standard, and the mixture was stirred. The brain sample was homogenized after addition of 2 ml water, an aliquot of the homogenate was taken and 3-fold amount of ethanol containing an internal standard
20 was added thereto and stirred. The cerebrospinal fluid was stirred after adding 3-fold amount of ethanol containing an internal standard. The samples thus obtained were allowed to stand at -20°C for 20 minutes, and then centrifuged (4°C, 12,000 g, 10 minutes). The supernatant was analyzed by LC/MS/MS, and
25 the concentration of the test compound in the plasma, brain, and cerebrospinal fluid were measured by the method using a relative calibration curve.

The results revealed that concentrations of the compound of Example 6 in the brain, cerebrospinal fluid and plasma were

1.10 nmol/g, 0.033 μ M and 2.77 μ M respectively, 2 hours after oral administration (10 mg/kg).

The compounds of the formula (I) can be administered orally or parenterally and, by formulating into a suitable administrable form, may be administered as a therapeutic agent for various diseases, including, for example, cardiovascular disorders such as angina, acute or congestive heart failure, myocardial infarction, hypertension, nephropathy, electrolyte abnormality, vasospasm, arteriosclerosis, etc., central nervous system disorders such as bulimia, depression, anxiety, seizure, epilepsy, dementia, pain, alcoholism, drug withdrawal, circadian rhythm disorders, schizophrenia, memory impairment, sleep disorders, cognitive impairment, etc., metabolic diseases such as obesity, diabetes, hormone abnormality, hypercholesterolemia, hyperlipidemia, gout, fatty liver, etc., genital or reproductive disorders such as infertility, preterm labor, sexual dysfunction, etc., gastro-intestinal disorders, respiratory disorder, inflammatory diseases or glaucoma, and the like, also for example, atherosclerosis, hypogonadism, hyperandrogenism, polycystic ovary syndrome, hirsutism, gastro-intestinal motility disorder, obesity-related gastro-esophageal reflux, obesity hypoventilation (Pickwickian syndrome), sleep apnea, inflammation, systemic inflammation of the vasculature, osteoarthritis, insulin resistance, bronchoconstriction, alcohol preference, metabolic syndrome, Alzheimer's disease, cardiac hypertrophy, left ventricular hypertrophy, hypertriglyceridemia, low HDL cholesterol, cardiovascular disorders such as coronary heart disease (CHD), cerebrovascular

disease, stroke, peripheral vascular disease, sudden death, gallbladder diseases, cancer (breast, endometrial, colon), breathlessness, hyperuricemia, impaired fertility, low back pain, or increased anesthetic risk, and the like. In clinical
5 use, the compounds of this invention may be administered after being formulated, together with pharmaceutically acceptable additives, into an appropriate preparation according to the mode of administration. As for said additives, those which are usually used in the field of pharmaceutical formulation may be
10 used, for example, gelatin, lactose, sucrose, titanium oxide, starch, crystalline cellulose, hydroxypropyl methylcellulose, carboxymethylcellulose, corn starch, microcrystalline wax, white petrolatum, magnesium methasilicate aluminate, anhydrous calcium phosphate, citric acid, trisodium citrate,
15 hydroxypropyl cellulose, sorbitol, sorbitan fatty acid ester, polysorbate, sucrose fatty acid ester, polyoxyethylene, hydrogenated castor oil, polyvinylpyrrolidone, magnesium stearate, light silicic anhydride, talc, vegetable oil, benzyl alcohol, gum arabic, propylene glycol, polyalkylene glycol,
20 cyclodextrin or hydroxypropyl cyclodextrin and the like.

A mixture with said additives may be formulated into the form of solid preparations (for example tablets, capsules, granules, powder, suppositories); or liquid preparations (for example syrups, elixirs, injections). Such preparations may be
25 formulated according to the techniques well-known in the art of pharmaceutical formulation. Liquid preparations may be in the form of preparations which are dissolved or suspended in water or other appropriate media when used, and injectable preparations in particular may be dissolved or suspended in

physiological saline or glucose solution if necessary, optionally together with a buffer and a preservative.

When compounds of this invention are used clinically, for example, a daily dose for an adult is 0.01-100 mg/kg, preferably 5 0.03-1 mg/kg with simultaneous or divided administration when administered orally, and 0.001-10 mg/kg, preferably 0.001-0.1 mg/kg, more preferably 0.01-0.1 mg/kg with simultaneous or divided administration when administered parenterally, though the dose and the frequency of dosage may vary depending upon 10 the sex, age, body weight, the degree of symptoms and the kind and range of the desired treatment effects.

An ordinarily skilled physician, veterinarian or clinician can readily determine and prescribe the effective amount of the drug required to prevent, suppress or arrest the 15 progress of diseases.

All the said preparations may contain 1.0 to 100 wt.%, preferably 1.0 to 60 wt.% of compounds of this invention and may also contain other therapeutically effective compounds.

The compounds of the present invention can be used in 20 combination with other agents useful for treating metabolic disorders and/or eating disorders. The individual component of such combinations can be administered separately at different times or concurrently in divided or single combination forms during the course of therapy. The instant 25 invention is therefore to be understood as embracing all such regimes of simultaneous or divided administration and the term "administering" is to be interpreted accordingly. The scope of combinations of the compounds of this invention with other agents useful for treating metabolic disorders and/or eating

disorders includes in principle any combination of any pharmaceutical composition useful for treating metabolic disorders and/or eating disorders.

Diabetes is caused by multiple factors and is most simply
5 characterized by elevated levels of plasma glucose (hyperglycemia) in the fasting state. There are two generally recognized forms of diabetes: type 1 diabetes, or insulin-dependent diabetes mellitus (IDDM), in which patients produce little or no insulin, the hormone which regulates
10 glucose utilization, and type 2 diabetes, or noninsulin-dependent diabetes mellitus (NIDDM), wherein patients produce insulin and even exhibit hyperinsulinemia (plasma insulin levels that are the same or even elevated in comparison with non-diabetic subjects), while at the same time
15 demonstrating hyperglycemia. Type 1 diabetes is typically treated with exogenous insulin administered via injection. However, type 2 diabetics often develop "insulin resistance", such that the effect of insulin in stimulating glucose and lipid metabolism in the main insulin-sensitive tissues, namely,
20 muscle, liver and adipose tissues, is diminished. Patients who are insulin resistant but not diabetic have elevated insulin levels that compensate for their insulin resistance, so that serum glucose levels are not elevated. In patients with NIDDM, the plasma insulin levels, even when they are elevated, are
25 insufficient to overcome the pronounced insulin resistance, resulting in hyperglycemia.

Insulin resistance is primarily due to a receptor binding defect that is not yet completely understood. Resistance to insulin results in insufficient activation of glucose uptake,

diminished oxidation of glucose and storage of glycogen in muscle, inadequate insulin repression of lipolysis in adipose tissue and inadequate glucose production and secretion by the liver.

5 The persistent or uncontrolled hyperglycemia that occurs in diabetics is associated with increased morbidity and premature mortality. Type 2 diabetics are at increased risk of developing cardiovascular complications, e.g., atherosclerosis, coronary heart disease, stroke, peripheral
10 vascular disease, hypertension, nephropathy, neuropathy and retinopathy.

Non-insulin dependent diabetes is also associated with cardiac hypertrophy, in particular left ventricular hypertrophy (Devereux, R. B., Circulation, 101:2271-2276
15 (2000)). Cardiac hypertrophy, such as left ventricular hypertrophy, is due to the response of the heart to chronic pressure or volume overload. Left ventricular hypertrophy (LVH) is characterized by thickening of the left ventricular wall, including increased left ventricular mass and increased
20 left ventricular wall thickness, and is defined as a left ventricular mass index exceeding 131 g/m^2 of the body surface area in men, and 100 g/m^2 in women (Savage et al., The Framingham Study, Circulation, 75 (1 Pt 2): 26-33 (1987)).

Left ventricular hypertrophy is independently associated
25 with increased incidence of cardiovascular disease, such as congestive heart failure, ischaemic heart disease, cardiovascular and all-cause mortality, sudden death, and stroke. Regression of left ventricular hypertrophy has been associated with a reduction in cardiovascular risk. It has also

been found that the incidence of morbid events in patients with progression of left ventricular hypertrophy is greater than in patients with regression of left ventricular hypertrophy.

Current treatments for hypertrophy include
5 non-pharmacological interventions, such as weight reduction, sodium restriction, and aerobic physical exercise can reduce left ventricular mass (Ghali, J.K. et al., American Journal of Geriatric Cardiology, 6:38-49 (1997)).

Many patients who have insulin resistance but have not
10 yet developed type 2 diabetes are also at a risk of developing metabolic syndrome, also referred to as syndrome X, insulin resistance syndrome, or plurimetabolic syndrome. The period of 5 to 10 years preceding the development of impaired glucose tolerance is associated with a number of hormonal imbalances,
15 which give rise to an enlargement of visceral fat mass, hypertension, insulin resistance, and hyperlipidemia (Bjornstop, P., Current Topics in Diabetes Research, eds. Belfore, F., Bergman, R. N., and Molinath, G. M., Front Diabetes, Basel, Karger, 12:182-192 (1993)). Similarly, metabolic
20 syndrome is characterized by insulin resistance, along with abdominal obesity, hyperinsulinemia, high blood pressure, low HDL and high VLDL. Although the causal relationship between the various components of metabolic syndrome remains to be confirmed, insulin resistance appears to play an important role
25 (Requen, G.M., et al., N. Eng. J. Med. 334:374-381 (1996); Despres, J-P., et al., N. Engl. J. Med. 334:952-957 (1996); Wajchenberg, B. L., et al., Diabetes /Metabolism Rev. 10:19-29 (1994)). Metabolic syndrome patients, whether or not they develop overt diabetes mellitus, are at increased risk of

developing the cardiovascular complications listed above. Associations have also been found between left ventricular hypertrophy and metabolic syndrome (Marcus, R. et al. Circulation, 90:928-936 (1994); Lind, L. et al., J Hypertens. 5 13:433-38 (1995); Paolisso, G et al., Am J Hypertens., 10:1250-1256 (1997)).

Diabetes is treated with a variety of therapeutic agents including insulin sensitizers, such as PPAR γ agonists, such as glitazones; biguanides; protein tyrosine phosphatase-1B 10 inhibitors; dipeptidyl peptidase IV inhibitors; insulin; insulin mimetics; sulfonylureas; meglitinides; α -glucoside hydrolase inhibitors; and α -amylase inhibitors.

Increasing the plasma level of insulin by administration of sulfonylureas (e.g. tolbutamide and glipizide) or 15 meglitinides, which stimulate the pancreatic β -cells to secrete more insulin, and/or by injection of insulin when sulfonylureas or meglitinides become ineffective, can result in insulin concentrations high enough to stimulate insulin-resistant tissues. However, dangerously low levels of plasma glucose can 20 result, and increasing insulin resistance due to the even higher plasma insulin levels can occur. The biguanides increase insulin sensitivity resulting in some correction of hyperglycemia. Metformin monotherapy is often used for treating type 2 diabetic patients who are also obese and/or 25 dyslipidemic. Lack of appropriate response to metformin is often followed by treatment with sulfonylureas, thiazolidinediones, insulin, or alpha glucosidase inhibitors. However, the two biguanides, phenformin and metformin, can also induce lactic acidosis and nausea/diarrhea, respectively.

Alpha glucosidase inhibitors, such as acarbose, work by delaying absorption of glucose in the intestine. Alpha-amylase inhibitors inhibit the enzymatic degradation of starch or glycogen into maltose, which also reduces the amounts of bioavailable sugars.

The glitazones, also known as thiazolidinediones (i.e. 5-benzylthiazolidine-2,4-diones), are a more recently described class of compounds with potential for a novel mode of action in ameliorating many symptoms of type 2 diabetes. These agents substantially increase insulin sensitivity in muscle, liver and adipose tissue in several animal models of type 2 diabetes resulting in partial or complete correction of the elevated plasma levels of glucose without occurrence of hypoglycemia. The glitazones that are currently marketed are agonists of the peroxisome proliferator activated receptor (PPAR) gamma subtype. PPAR-gamma agonism is generally believed to be responsible for the improved insulin sensitization that is observed with the glitazones. Newer PPAR agonists that are being developed for treatment of Type 2 diabetes and/or dyslipidemia are agonists of one or more of the PPAR alpha, gamma and delta subtypes.

However, treatment of diabetes with PPAR γ agonists has been associated with cardiac hypertrophy, or an increase in heart weight. Recent labeling revisions for Avandia (rosiglitazone maleate), a PPAR γ agonist, indicate that patients may experience fluid accumulation and volume-related events such as edema and congestive heart failure. Cardiac hypertrophy related to PPAR γ agonist treatment is typically treated by withdrawing PPAR treatment.

Treatment of type 2 diabetes also typically includes physical exercise, weight control and dieting. While physical exercise and reductions in dietary intake of calories will dramatically improve the diabetic condition, compliance with
5 this treatment is very poor because of well-entrenched sedentary lifestyles and excess food consumption, especially of foods containing high amounts of saturated fat. However, weight reduction and increased exercise are difficult for most people with diabetes.

10 Abnormal glucose homeostasis is also associated both directly and indirectly with obesity, hypertension and alterations in lipid, lipoprotein and apolipoprotein metabolism. Obesity increases the likelihood of insulin resistance, and increases the likelihood that the resulting
15 insulin resistance will increase with increasing body weight. Therefore, therapeutic control of glucose homeostasis, lipid metabolism, obesity and hypertension are critically important in the clinical management and treatment of diabetes mellitus.

Obesity, which can be defined as a body weight more than
20 20% above the ideal body weight, is a major health concern in Western societies. It is estimated that about 97 million adults in the United States are overweight or obese. Obesity is the result of a positive energy balance, as a consequence of increased ratio of caloric intake to energy expenditure. The
25 molecular factors regulating food intake and body weight balance are incompletely understood. [B. Staels et al., J. Biol. Chem. 270(27), 15958 (1995); F. Lonnquist et al., Nature Medicine 1(9), 950 (1995)]. Although the genetic and/or

environmental factors leading to obesity are poorly understood, several genetic factors have been identified.

Epidemiological studies have shown that increasing degrees of overweight and obesity are important predictors of decreased life expectancy. Obesity causes or exacerbates many health problems, both independently and in association with other diseases. The medical problems associated with obesity, which can be serious and life-threatening, include type 2 diabetes mellitus, hypertension, elevated plasma insulin concentrations, insulin resistance, dyslipidemias, hyperlipidemia, endometrial, breast, prostate, kidney and colon cancer, osteoarthritis; respiratory complications, such as obstructive sleep apnea, gallstones, arteriosclerosis, heart disease, abnormal heart rhythms, and heart arrhythmias (Kopelman, P.G., Nature 404, 635-643 (2000)). Obesity is also associated with metabolic syndrome, cardiac hypertrophy, in particular left ventricular hypertrophy, premature death, and with a significant increase in mortality and morbidity from stroke, myocardial infarction, congestive heart failure, coronary heart disease, and sudden death.

Abdominal obesity has been linked with a much higher risk of coronary artery disease, and with three of its major risk factors: high blood pressure, diabetes that starts in adulthood, and high levels of fats (lipids) in the blood. Losing weight dramatically reduces these risks. Abdominal obesity is further closely associated with glucose intolerance, hyperinsulinemia, hypertriglyceridemia, and other disorders associated with metabolic syndrome (syndrome X), such as raised high blood pressure, decreased levels of high density

lipoproteins (HDL) and increased levels of very low density lipoproteins (VLDL) (Montague et al., Diabetes, 2000, 49: 883-888).

Obesity and obesity-related disorders, such as diabetes, are often treated by encouraging patients to lose weight by reducing their food intake or by increasing their exercise level, thereby increasing their energy output. A sustained weight loss of 5% to 10% of body weight has been shown to improve the comorbidities associated with obesity, such as diabetes, and can lead to improvement of obesity-related disorders such as diabetes, left ventricular hypertrophy, osteoarthritis, and pulmonary and cardiac dysfunction.

Weight loss drugs used for the treatment of obesity include orlistat (Davidson, M.H. et al. (1999) JAMA 281:235-42), dexfenfluramine (Guy Grand, B. et al. (1989) Lancet 2:1142-5), sibutramine (Bray, G. A. et al. (1999) Obes. Res. &:189-98) and phentermine (Douglas, A. et al. (1983) Int. J. Obes. 7:591-5). However, the side effects of these drugs and anti-obesity agents may limit their use. Dexfenfluramine was withdrawn from the market because of suspected heart valvulopathy; orlistat is limited by gastrointestinal side effects; and the use of sibutramine is limited by its cardiovascular side effects which have led to reports of deaths and its withdrawal from the market in Italy.

The term "diabetes," as used herein, includes both insulin-dependent diabetes mellitus (i.e., IDDM, also known as type 1 diabetes) and non-insulin-dependent diabetes mellitus (i.e., NIDDM, also known as Type 2 diabetes). Type 1 diabetes, or insulin-dependent diabetes, is the result of an absolute

deficiency of insulin, the hormone which regulates glucose utilization. Type 2 diabetes, or insulin-independent diabetes (i.e., non-insulin-dependent diabetes mellitus), often occurs in the face of normal, or even elevated levels of insulin and
5 appears to be the result of the inability of tissues to respond appropriately to insulin. Most of the Type 2 diabetics are also obese. The compositions of the present invention are useful for treating both Type 1 and Type 2 diabetes. The compositions are especially effective for treating Type 2 diabetes. The
10 compositions of the present invention are also useful for treating and/or preventing gestational diabetes mellitus.

Treatment of diabetes mellitus refers to the administration of a compound or combination of the present invention to treat diabetes. One outcome of treatment may be
15 decreasing the glucose level in a subject with elevated glucose levels. Another outcome of treatment may be decreasing insulin levels in a subject with elevated insulin levels. Another outcome of treatment is decreasing plasma triglycerides in a subject with elevated plasma triglycerides. Another outcome
20 of treatment is decreasing LDL cholesterol in a subject with high LDL cholesterol levels. Another outcome of treatment is increasing HDL cholesterol in a subject with low HDL cholesterol levels. Another outcome of treatment is increasing insulin sensitivity. Another outcome of treatment may be enhancing
25 glucose tolerance in a subject with glucose intolerance. Yet another outcome of treatment may be decreasing insulin resistance in a subject with increased insulin resistance or elevated levels of insulin.

Prevention of diabetes mellitus refers to the administration of a compound or combination of the present invention to prevent the onset of diabetes in a subject in need thereof.

5 The term "hypertension" as used herein includes essential, or primary, hypertension wherein the cause is not known or where hypertension is due to greater than one cause, such as changes in both the heart and blood vessels; and secondary hypertension wherein the cause is known. Causes of secondary hypertension
10 include, but are not limited to obesity; kidney disease; hormonal disorders; use of certain drugs, such as oral contraceptives, corticosteroids, cyclosporin, and the like. The term "hypertension" encompasses high blood pressure, in which both the systolic and diastolic pressure levels are
15 elevated, and isolated systolic hypertension, in which only the systolic pressure is elevated to greater than or equal to 140 mm Hg, while the diastolic pressure is less than 90 mm Hg. One outcome of treatment is decreasing blood pressure in a subject with high blood pressure.

20 Dyslipidemias or disorders of lipid metabolism, include various conditions characterized by abnormal concentrations of one or more lipids (i.e. cholesterol and triglycerides), and/or apolipoproteins (i.e., apolipoproteins A, B, C and E), and/or lipoproteins (i.e., the macromolecular complexes formed by the
25 lipid and the apolipoprotein that allow lipids to circulate in blood, such as LDL, VLDL and IDL). Hyperlipidemia is associated with abnormally high levels of lipids, LDL and VLDL cholesterol, and/or triglycerides.

The term "metabolic syndrome", also known as syndrome X, is defined in the Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (ATP-III). E.S. Ford et al., JAMA, vol. 287 (3), Jan. 16, 2002, pp 356-359. Briefly, a person is defined as having metabolic syndrome if the person has three or more of the following symptoms: abdominal obesity, hypertriglyceridemia, low HDL cholesterol, high blood pressure, and high fasting plasma glucose. The criteria for these are defined in ATP-III.

The term "left ventricular hypertrophy" (LVH) as used herein includes three patterns of left ventricular hypertrophy that have been identified based on left ventricular mass index (LVMI = left ventricular mass in grams divided by body surface area in meters²) and relative wall thickness (RWT = 2 x posterior wall thickness/left ventricular end diastolic diameter). Concentric LVH is typically exemplified by a left ventricular mass index of 144 and a relative wall thickness of 0.52; eccentric LVH is typically exemplified by a left ventricular mass index of 136 and a relative wall thickness of 0.38; and concentric left ventricular remodeling which is typically exemplified by a LVMI of 93 and a relative wall thickness of 0.49. Normal LVMI are typically 85 and normal RWT approximately 0.36. Patients with concentric left ventricular (LV) remodeling have a cardiovascular risk intermediate between those with normal left ventricular structure and those with left ventricular hypertrophy.

One outcome of treatment of diabetes while minimizing cardiac hypertrophy, or left ventricular hypertrophy, may be

a decrease in ventricular mass. Another outcome of treatment of diabetes while minimizing cardiac hypertrophy or left ventricular hypertrophy may be a decrease in the rate of increase of ventricular mass. Another outcome of treatment of diabetes while minimizing cardiac hypertrophy or left ventricular hypertrophy may be a decrease in ventricular wall thickness. Another outcome of treatment of diabetes while minimizing cardiac hypertrophy of left ventricular hypertrophy may be the decrease in the rate of increase in ventricular wall thickness.

The term "obesity" as used herein is a condition in which there is an excess of body fat. The operational definition of obesity is based on the Body Mass Index (BMI), which is calculated as body weight per height in meters squared (kg/m^2). "Obesity" refers to a condition whereby an otherwise healthy subject has a Body Mass Index (BMI) greater than or equal to $30 \text{ kg}/\text{m}^2$, or a condition whereby a subject with at least one co-morbidity has a BMI greater than or equal to $27 \text{ kg}/\text{m}^2$. An "obese subject" is an otherwise healthy subject with a Body Mass Index (BMI) greater than or equal to $30 \text{ kg}/\text{m}^2$ or a subject with at least one co-morbidity with a BMI greater than or equal to $27 \text{ kg}/\text{m}^2$. A "subject at risk of obesity" is an otherwise healthy subject with a BMI of $25 \text{ kg}/\text{m}^2$ to less than $30 \text{ kg}/\text{m}^2$ or a subject with at least one co-morbidity with a BMI of $25 \text{ kg}/\text{m}^2$ to less than $27 \text{ kg}/\text{m}^2$.

The increased risks associated with obesity occur at a lower Body Mass Index (BMI) in Asians. In Asian countries, including Japan, "obesity" refers to a condition whereby a subject with at least one obesity-induced or obesity-related

co-morbidity, that requires weight reduction or that would be improved by weight reduction, has a BMI greater than or equal to 25 kg/m². In Asian countries, including Japan, an "obese subject" refers to a subject with at least one obesity-induced or obesity-related co-morbidity that requires weight reduction or that would be improved by weight reduction, with a BMI greater than or equal to 25 kg/m². In Asia-Pacific, a "subject at risk of obesity" is a subject with a BMI of greater than 23 kg/m² to less than 25 kg/m².

As used herein, the term "obesity" is meant to encompass all of the above definitions of obesity.

Obesity-induced or obesity-related co-morbidities include, but are not limited to, diabetes, non-insulin dependent diabetes mellitus - type 2, diabetes associated with obesity, impaired glucose tolerance, impaired fasting glucose, insulin resistance syndrome, dyslipidemia, hypertension, hypertension associated with obesity, hyperuricacidemia, gout, coronary artery disease, myocardial infarction, angina pectoris, sleep apnea syndrome, Pickwickian syndrome, fatty liver; cerebral infarction, cerebral thrombosis, transient ischemic attack, orthopedic disorders, arthritis deformans, lumbodynia, emmeniopathy, and infertility. In particular, co-morbidities include: hypertension, hyperlipidemia, dyslipidemia, glucose intolerance, cardiovascular disease, sleep apnea, diabetes mellitus, and other obesity-related conditions.

Treatment of obesity and obesity-related disorders refers to the administration of the compounds or combinations of the present invention to reduce or maintain the body weight

of an obese subject. One outcome of treatment may be reducing the body weight of an obese subject relative to that subject's body weight immediately before the administration of the compounds or combinations of the present invention. Another outcome of treatment may be preventing body weight regain of body weight previously lost as a result of diet, exercise, or pharmacotherapy. Another outcome of treatment may be decreasing the occurrence of and/or the severity of obesity-related diseases. The treatment may suitably result in a reduction in food or calorie intake by the subject, including a reduction in total food intake, or a reduction of intake of specific components of the diet such as carbohydrates or fats; and/or the inhibition of nutrient absorption; and/or the inhibition of the reduction of metabolic rate; and in weight reduction in patients in need thereof. The treatment may also result in an alteration of metabolic rate, such as an increase in metabolic rate, rather than or in addition to an inhibition of the reduction of metabolic rate; and/or in minimization of the metabolic resistance that normally results from weight loss.

Prevention of obesity and obesity-related disorders refers to the administration of the compounds or combinations of the present invention to reduce or maintain the body weight of a subject at risk of obesity. One outcome of prevention may be reducing the body weight of a subject at risk of obesity relative to that subject's body weight immediately before the administration of the compounds or combinations of the present invention. Another outcome of prevention may be preventing body weight regain of body weight previously lost as a result

of diet, exercise, or pharmacotherapy. Another outcome of prevention may be preventing obesity from occurring if the treatment is administered prior to the onset of obesity in a subject at risk of obesity. Another outcome of prevention may be decreasing the occurrence and/or severity of obesity-related disorders if the treatment is administered prior to the onset of obesity in a subject at risk of obesity. Moreover, if treatment is commenced in already obese subjects, such treatment may prevent the occurrence, progression or severity of obesity-related disorders, such as, but not limited to, arteriosclerosis, Type 2 diabetes, polycystic ovary disease, cardiovascular diseases, osteoarthritis, dermatological disorders, hypertension, insulin resistance, hypercholesterolemia, hypertriglyceridemia, and cholelithiasis.

The term "atherosclerosis" as used herein encompasses vascular diseases and conditions that are recognized and understood by physicians practicing in the relevant fields of medicine. Atherosclerotic cardiovascular disease, coronary heart disease (also known as coronary artery disease or ischemic heart disease), cerebrovascular disease and peripheral vessel disease are all clinical manifestations of atherosclerosis and are therefore encompassed by the terms "atherosclerosis" and "atherosclerotic disease." The combination comprised of a therapeutically effective amount of an anti-obesity agent in combination with a therapeutically effective amount of an anti-diabetic agent may be administered to prevent or reduce the risk of occurrence, or recurrence where the potential exists, of a coronary heart disease event, a cerebrovascular event, or

intermittent claudication. Coronary heart disease events are intended to include CHD death, myocardial infarction (i.e., a heart attack), and coronary revascularization procedures. Cerebrovascular events are intended to include ischemic or hemorrhagic stroke (also known as cerebrovascular accidents) and transient ischemic attacks. Intermittent claudication is a clinical manifestation of peripheral vessel disease. The term "atherosclerotic disease event" as used herein is intended to encompass coronary heart disease events, cerebrovascular events, and intermittent claudication. It is intended that persons who have previously experienced one or more non-fatal atherosclerotic disease events are those for whom the potential for recurrence of such an event exists.

Circadian rhythms affect a variety of physiological parameters: rest-activity, sleep-wake cycles, body temperature, rhythms in hormone levels, oscillations in general physiology and the like. When these parameters are out of synchrony with the daily clock, a circadian rhythm imbalance occurs which can affect physiology, performance on a variety of tasks and one's emotional well being. The present invention is useful, for example, in the prevention or treatment of conditions associated with circadian rhythmicity as well as mental and physical disorders associated with travel across time zones and with rotating shift-work schedules.

In another embodiment, the present invention provides a method for the prevention or treatment of a circadian rhythm disorder in a mammal, including time-zone change (jet-lag) syndrome, shift-work sleep disorder, delayed sleep-phase syndrome, advanced sleep-phase syndrome, and non-24-hour

sleep-wake disorder, which comprises administering to the mammal an effective amount of a NPY Y5 receptor antagonist.

In another embodiment, the present invention provides a method for shortening the time of re-entrainment (return to
5 normal entrainment of the circadian rhythms; synchronized to the environmental light-dark cycle) in a subject following a shift in the sleep-wake cycle which comprises administering to the subject an appropriate amount of a NPY Y5 antagonist.

In another embodiment, the present invention provides a
10 method for alleviating the effects of jet lag in a traveler, especially a mammal, which comprises administering to the traveler an alertness increasing amount of a NPY Y5 antagonist. The purpose of this embodiment is to assist the body to adjust physiologically to the changes in sleep and feeding patterns
15 when crossing several time zones.

In another more preferred embodiment, the present invention provides a method for resetting the internal circadian clock in a subject to match the subject's current activity/sleep cycle. For example shift workers changing from
20 a day to a night shift or vice versa, which comprises administering to the subject an appropriate amount of a NPY Y5 antagonist.

The present invention is further directed to the use of NPY Y5 antagonist, for enhancing or improving sleep quality as
25 well as preventing and treating sleep disorders and sleep disturbances in a mammal. In particular, the present invention provides a method for enhancing or improving sleep quality by increasing sleep efficiency and augmenting sleep maintenance. In addition, the present invention provides a method for

preventing and treating sleep disorders and sleep disturbances in a mammal which comprising the administration of a NPY Y5 antagonist. The present invention further provides a pharmaceutical composition for enhancing or improving sleep
5 quality and increasing sleep efficiency and sleep maintenance. The present invention is useful for the treatment of sleep disorders, including Disorders of Initiating and Maintaining Sleep (insomnias) ("DIMS") which can arise from psychophysiological causes, as a consequence of psychiatric
10 disorders (particularly related to anxiety), from drugs and alcohol use and abuse (particularly during withdrawal stages), childhood onset DIMS, nocturnal myoclonus and restless legs and non specific REM disturbances as seen in ageing.

The following outcomes in a subject which are provided
15 by the present invention may be correlated to enhancement in sleep quality: an increase in the value which is calculated from the time that a subject sleeps divided by the time that a subject is attempting to sleep; a decrease in sleep latency (the time it takes to fall asleep); a decrease in the number
20 of awakenings during sleep; a decrease in the time spent awake following the initial onset of sleep; an increase in the total amount of sleep; an increase the amount and percentage of REM sleep; an increase in the duration and occurrence of REM sleep; a reduction in the fragmentation of REM sleep; an increase in
25 the amount and percentage of slow-wave (i.e. stage 3 or 4) sleep; an increase in the amount and percentage of stage 2 sleep; a decrease in the number of awakenings, especially in the early morning; an increase in daytime alertness; and increased sleep maintenance. Secondary outcomes which may be provided by the

present invention include enhanced cognitive function and increased memory retention. A "method for enhancing the quality of sleep" refers to a method that results in outcomes in a subject which may be correlated to enhancement in sleep
5 quality, including, but not limited to, the outcomes correlated to enhancement of sleep quality as defined above.

The present invention is further useful for the prevention and treatment of sleep disorders and sleep disturbances including sleep problems associated with insomnia,
10 hypersomnia, sleep apnea, narcolepsy, nocturnal myoclonus, REM sleep interruptions, jet-lag, shift workers' sleep disturbances, dysomnias, night terror, night eating syndrome, insomnias associated with depression or with emotional/mood disorders, dysfunctions associated with sleep (parasomnias),
15 as well as sleep walking and enuresis, as well as sleep disorders which accompany aging. Sleep disorders and sleep disturbances are generally characterized by difficulty in initiating or maintaining sleep or in obtaining restful or enough sleep.

In addition, certain drugs may also cause reductions in
20 REM sleep as a side effect and the present invention may be used to correct those types of sleeping disorders as well. The present invention would also be of benefit in the treatment of syndromes such as fibromyalgia which are manifested by non-restorative sleep and muscle pain or sleep apnea which is
25 associated with respiratory disturbances during sleep. It will be clear to one skilled in the art that the present invention is not limited to just sleep disorders and sleep disturbances, but is applicable to a wide variety of conditions which result from a diminished quality of sleep.

The present invention is also concerned with treatment and prevention of these conditions, and with the use of a NPY Y5 antagonist, combinations, and compositions thereof, for the manufacture of a medicament useful for treating or preventing these conditions.

In the present invention, it is preferred that the subject mammal is a human. Although the present invention is applicable both old and young people, it may find greater application in elderly people. Further, although the invention may be employed to enhance the sleep of healthy people, it may be especially beneficial for enhancing the sleep quality of people suffering from sleep disorders or sleep disturbances.

The compositions of the present invention may be used in combination with other drugs that may also be useful in the treatment, prevention, or control of disorders, such as hypertension, hypertension associated with obesity, hypertension-related disorders, cardiac hypertrophy, left ventricular hypertrophy, and metabolic syndrome, obesity and obesity-related disorders, for which compounds comprising the compositions are useful. Such other drugs may be administered, by a route and in an amount commonly used therefore, contemporaneously or sequentially with a composition of the present invention. When a composition of the present invention is used contemporaneously with one or more other drugs, a pharmaceutical composition in unit dosage form containing such other drugs and the composition of the present invention is preferred. However, the combination therapy also includes therapies in which the composition of the present invention and one or more other drugs are administered on different

overlapping schedules. It is also contemplated that when used in combination with one or more other active ingredients, the composition of the present invention and the other active ingredients may be used in lower doses than when each is used
5 singly. Accordingly, the pharmaceutical compositions of the present invention include those that contain one or more other active ingredients, in addition to a composition of the present invention.

Examples of other active ingredients that may be
10 administered in combination with a composition of the present invention, and either administered separately or in the same pharmaceutical composition, include, but are not limited to:

(a) anti-diabetic agents such as (i) PPAR γ agonists such as glitazones (e.g. ciglitazone; darglitazone; englitazone; isaglitazone (MCC-555); pioglitazone; rosiglitazone; troglitazone; BRL49653; CLX-0921; 5-BTZD, and the like), and GW-0207, LG-100641, and LY-300512, and the like; (ii) biguanides such as buformin; metformin; and phenformin, and the like; (iii) protein tyrosine phosphatase-1B (PTP-1B)
15 inhibitors; (iv) sulfonylureas such as acetohexamide; chlorpropamide; diabinese; glibenclamide; glipizide; glyburide; glimepiride; gliclazide; glipentide; gliquidone; glisolamide; tolazamide; and tolbutamide, and the like; (v) meglitinides such as repaglinide, and nateglinide, and the
20 like; (vi) alpha glucoside hydrolase inhibitors such as acarbose; adiposine; camiglibose; emiglitate; miglitol; voglibose; pradiamicin-Q; salbostatin; CKD-711; MDL-25,637; MDL-73,945; and MOR 14, and the like; (vii) alpha-amylase inhibitors such as tendamistat, trestatin, and A1-3688, and the

like; (viii) insulin secretagogues such as linoglriride; and A-4166, and the like; (ix) fatty acid oxidation inhibitors, such as clomoxir, and etomoxir, and the like; (x) A2 antagonists, such as midaglizole; isaglidole; deriglidole; idazoxan; 5 earoxan; and fluparoxan, and the like; (xi) insulin or insulin mimetics, such as biota, LP-100, novarapid, insulin detemir, insulin lispro, insulin glargine, insulin zinc suspension (lente and ultralente); Lys-Pro insulin, GLP-1 (73-7) (insulintropin); and GLP-1 (7-36)-NH₂, and the like; (xii) 10 non-thiazolidinediones such as JT-501, and farglitazar (GW-2570/GI-262579), and the like; (xiii) PPAR α / γ dual agonists such as MK-0767, CLX-0940, GW-1536, GW1929, GW-2433, KRP-297, L-796449, LR-90, and SB 219994, and the like; (xiv) other insulin sensitizing drugs; and (xv) VPAC2 receptor agonists;

15 (b) lipid lowering agents such as (i) bile acid sequestrants such as, cholestyramine, colesevelem, colestipol, dialkylaminoalkyl derivatives of a cross-linked dextran; Colestid®; LoCholest®; and Questran®, and the like; (ii) HMG-CoA reductase inhibitors such as atorvastatin, 20 itavastatin, fluvastatin, lovastatin, pravastatin, rivastatin, rosuvastatin, simvastatin, and ZD-4522, and the like; (iii) HMG-CoA synthase inhibitors; (iv) cholesterol absorption inhibitors such as stanol esters, beta-sitosterol, sterol glycosides such as tiqueside; and azetidinones such as 25 ezetimibe, and the like; (v) acyl coenzyme A -cholesterol acyl transferase (ACAT) inhibitors such as avasimibe, eflucimibe, KY505, SMP 797, and the like; (vi) CETP inhibitors such as JTT 705, torcetrapib, CP 532,632, BAY63-2149, SC 591, SC 795, and the like; (vii) squalene synthetase inhibitors; (viii)

anti-oxidants such as probucol, and the like; (ix) PPAR α agonists such as beclofibrate, benzaifibrate, ciprofibrate, clofibrate, etofibrate, fenofibrate, gemcabene, and gemfibrozil, GW 7647, BM 170744, LY518674; and other fibric acid derivatives, such as Atromid®, Lopid® and Tricor®, and the like; (x) FXR receptor modulators such as GW 4064, SR 103912, and the like; (xi) LXR receptor such as GW 3965, T9013137, and XTC0179628, and the like; (xii) lipoprotein synthesis inhibitors such as niacin; (xiii) renin angiotensin system inhibitors; (xiv) PPAR δ partial agonists; (xv) bile acid reabsorption inhibitors, such as BARI 1453, SC435, PHA384640, S8921, AZD7706, and the like; (xvi) PPAR δ agonists such as GW 501516, and GW 590735, and the like; (xvii) triglyceride synthesis inhibitors; (xviii) microsomal triglyceride transport (MTTP) inhibitors, such as inplitapide, LAB687, and CP346086, and the like; (xix) transcription modulators; (xx) squalene epoxidase inhibitors; (xxi) low density lipoprotein (LDL) receptor inducers; (xxii) platelet aggregation inhibitors; (xxiii) 5-LO or FLAP inhibitors; and (xiv) niacin receptor agonists; and

(c) anti-hypertensive agents such as (i) diuretics, such as thiazides, including chlorthalidone, chlorthiazide, dichlorophenamide, hydroflumethiazide, indapamide, and hydrochlorothiazide; loop diuretics, such as bumetanide, ethacrynic acid, furosemide, and torsemide; potassium sparing agents, such as amiloride, and triamterene; and aldosterone antagonists, such as spironolactone, epi renone, and the like; (ii) beta-adrenergic blockers such as acebutolol, atenolol, betaxolol, bevantolol, bisoprolol, bopindolol, carteolol,

carvedilol, celiprolol, esmolol, indenolol, metoprolol, nadolol, nebivolol, penbutolol, pindolol, propanolol, sotalol, tertatolol, tilisolol, and timolol, and the like; (iii) calcium channel blockers such as amlodipine, aranidipine, azelnidipine, barnidipine, benidipine, bepridil, cinaldipine, clevidipine, diltiazem, efonidipine, felodipine, gallopamil, isradipine, lacidipine, lemildipine, lercanidipine, nicardipine, nifedipine, nilvadipine, nimodepine, nisoldipine, nitrendipine, manidipine, pranidipine, and verapamil, and the like; (iv) angiotensin converting enzyme (ACE) inhibitors such as benazepril; captopril; cilazapril; delapril; enalapril; fosinopril; imidapril; losinopril; moexipril; quinapril; quinaprilat; ramipril; perindopril; perindropril; quanipril; spirapril; tenocapril; trandolapril, and zofenopril, and the like; (v) neutral endopeptidase inhibitors such as omapatrilat, cadoxatril and ecadotril, fosidotril, sampatrilat, AVE7688, ER4030, and the like; (vi) endothelin antagonists such as tezosentan, A308165, and YM62899, and the like; (vii) vasodilators such as hydralazine, clonidine, minoxidil, and nicotiny alcohol, and the like; (viii) angiotensin II receptor antagonists such as candesartan, eprosartan, irbesartan, losartan, prazosartan, tasosartan, telmisartan, valsartan, and EXP-3137, FI6828K, and RNH6270, and the like; (vix) α/β adrenergic blockers as nipradilol, arotinolol and amosulalol, and the like; (x) α 1 blockers, such as terazosin, urapidil, prazosin, bunazosin, trimazosin, doxazosin, naftopidil, indoramin, WHIP 164, and XEN010, and the like; (xi) α 2 agonists such as lofexidine, tiamenidine, moxonidine,

rilmidenidine and guanobenz, and the like; and (xii) aldosterone inhibitors, and the like; and

(d) anti-obesity agents, such as (i) 5HT (serotonin) transporter inhibitors, such as paroxetine, fluoxetine, 5 fenfluramine, fluvoxamine, sertraline, and imipramine; (ii) NE (norepinephrine) transporter inhibitors, such as GW 320659, despiramine, talsupram, and nomifensine; (iii) CB-1 (cannabinoid-1 receptor) antagonist/inverse agonists, such as rimonabant (Sanofi Synthelabo), SR-147778 (Sanofi Synthelabo), BAY 65-2520 (Bayer), and SLV 319 (Solvay), and 10 those disclosed in U.S. Patent Nos. 5,532,237, 4,973,587, 5,013,837, 5,081,122, 5,112,820, 5,292,736, 5,624,941 and US 6,028,084; and WO 96/33159, WO 98/33765, WO98/43636, WO98/43635, WO 01/09120, WO 01.96330, WO98/31227, WO98/41519, WO98/37061, 15 WO00/10967, WO00/10968, WO97/29079, WO99/02499, WO 01/58869, WO 02/076949, WO 01/64632, WO 01/64633, WO 01/64634, WO 03/006007, and WO 03/007887; and EPO Application No. EP-658546; (iv) ghrelin antagonists, such as those disclosed in WO 01/87335, and WO 02/08250; (v) H3 (histamine H3) antagonist/inverse 20 agonists, such as thioperamide, 3-(1H-imidazol-4-yl)propyl N-(4-pentenyl)carbamate, clobenpropit, iodophenpropit, imoproxifan, GT2394 (Gliatech), and A331440, and those disclosed in WO 02/15905; and O-[3-(1H-imidazol-4-yl)propanol]carbamates (Kiec-Kononowicz, 25 K. et al., Pharmazie, 55:349-55 (2000)), piperidine-containing histamine H3-receptor antagonists (Lazewska, D. et al., Pharmazie, 56:927-32 (2001)), benzophenone derivatives and related compounds (Sasse, A. et al., Arch. Pharm.(Weinheim) 334:45-52 (2001)), substituted N-phenylcarbamates

(Reidemeister, S. et al., Pharmazie, 55:83-6 (2000)), and proxifan derivatives (Sasse, A. et al., J. Med. Chem. 43:3335-43 (2000)); (vi) melanin-concentrating hormone 1 receptor (MCH1R) antagonists, such as T-226296 (Takeda),
5 SNP-7941 (Synaptic), and those disclosed in WO 01/82925, WO 01/87834, WO 02/051809, WO 02/06245, WO 02/076929, WO 02/076947, WO 02/04433, WO 02/51809, WO 02/083134, WO 02/094799, WO 03/004027, and Japanese Patent Application No. JP 13226269;
(vii) MCH2R (melanin concentrating hormone 2R)
10 agonist/antagonists; (viii) NPY1 (neuropeptide Y Y1) antagonists, such as BIBP3226, 2-[1-(5-chloro-3-isopropoxyloxycarbonylamino-phenyl)ethylamino]-6-[2-(5-ethyl-4-methyl-1,3-thiazol-2-yl)ethyl]-4-morpholinopyridine, BIBO 3304, LY-357897, CP-671906, and GI-264879A;
15 and those disclosed in U.S. Patent No. 6,001,836; and WO 96/14307, WO 01/23387, WO 99/51600, WO 01/85690, WO 01/85098, WO 01/85173, and WO 01/89528; (ix) NPY5 (neuropeptide Y Y5) antagonists, such as L-152,804, GW-569180A, GW-594884A, GW-587081X, GW-548118X; FR 235,208; FR226928, FR 240662,
20 FR252384; 1229U91, GI-264879A, CGP71683A, LY-377897, LY366377, PD-160170, SR-120562A, SR-120819A, JCF-104, and H409/22; and those compounds disclosed in U.S. Patent Nos. 6,140,354, 6,191,160, 6,258,837, 6,313,298, 6,337,332, 6,329,395, and 6,340,683; U.S. Patent Nos. 6,326,375; 6,329,395; 6,337,332;
25 6,335,345; European Patent Nos. EP-01010691, and EP-01044970; and PCT International Patent Publication Nos. WO 97/19682, WO 97/20820, WO 97/20821, WO 97/20822, WO 97/20823, WO 98/27063, WO 00/107409, WO 00/185714, WO 00/185730, WO 00/64880, WO 00/68197, WO 00/69849, WO 01/09120, WO 01/14376, WO 01/85714,

WO 01/85730, WO 01/07409, WO 01/02379, WO 01/02379, WO 01/23388,
WO 01/23389, WO 01/44201, WO 01/62737, WO 01/62738, WO 01/09120,
WO 02/20488, WO 02/22592, WO 02/48152, WO 02/49648 and WO
02/094789; and Norman et al., J. Med. Chem. 43:4288-4312 (2000);
5 (x) leptin, such as recombinant human leptin (PEG-OB, Hoffman
La Roche) and recombinant methionyl human leptin (Amgen); (xi)
leptin derivatives, such as those disclosed in Patent Nos.
5,552,524; 5,552,523; 5,552,522; 5,521,283; and PCT
International Publication Nos. WO 96/23513; WO 96/23514; WO
10 96/23515; WO 96/23516; WO 96/23517; WO 96/23518; WO 96/23519;
and WO 96/23520; (xii) opioid antagonists, such as nalmefene
(Revex®), 3-methoxynaltrexone, naloxone, and naltrexone; and
those disclosed in WO 00/21509; (xiii) orexin antagonists, such
as SB-334867-A; and those disclosed in WO 01/96302, WO 01/68609,
15 WO 02/51232, WO 02/51838, and WO 03/023561; (xiv) BRS3
(bombesin receptor subtype 3) agonists; (xv) CCK-A
(cholecystokinin-A) agonists, such as AR-R 15849, GI 181771,
JMV-180, A-71378, A-71623 and SR146131, and those disclosed in
US 5,739,106; (xvi) CNTF (ciliary neurotrophic factors), such
20 as GI-181771 (Glaxo-SmithKline); SR146131 (Sanofi
Synthelabo); butabindide; and PD170,292, PD 149164 (Pfizer);
(xvii) CNTF derivatives, such as axokine (Regeneron); and WO
94/09134, WO 98/22128, and WO 99/43813; (xviii) GHS (growth
hormone secretagogue receptor) agonists, such as NN703,
25 hexarelin, MK-0677, SM-130686, CP-424,391, L-692,429 and
L-163,255, and those disclosed in U.S. Patent No. 6358951, U.S.
Patent Application Nos. 2002/049196 and 2002/022637; and WO
01/56592, and WO 02/32888; (xix) 5HT2c (serotonin receptor 2c)
agonists, such as BVT933, DPCA37215, IK264; PNU 22394;

WAY161503, R-1065, and YM 348; and those disclosed in U.S. Patent No. 3,914,250; and WO 02/36596, WO 02/48124, WO 02/10169, WO 01/66548, WO 02/44152; WO 02/51844, WO 02/40456, and WO 02/40457; (xx) Mc3r (melanocortin 3 receptor) agonists; 5 (xxi) Mc4r (melanocortin 4 receptor) agonists, such as CHIR86036 (Chiron); ME-10142, and ME-10145 (Melacure), and those disclosed in WO 99/64002, WO 00/74679, WO 01/991752, WO 01/74844, WO 01/70708, WO 01/70337, WO 01/91752, WO 02/059095, WO 02/059107, WO 02/059108, WO 02/059117, WO 02/12166, WO 10 02/11715, WO 02/12178, WO 02/15909, WO 02/068387, WO 02/068388, WO 02/067869, WO 03/007949, and WO 03/009847; (xxii) monoamine reuptake inhibitors, such as sibutramine (Meridia ®/Reductil®) and a salt thereof, and those compounds disclosed in U.S. Patent Nos. 4,746,680, 4,806,570, and 5,436,272, and 15 U.S. Patent Publication No. 2002/0006964, and WO 01/27068, and WO 01/62341; (xxiii) serotonin reuptake inhibitors, such as dexfenfluramine, fluoxetine, and those in U.S. Patent No. 6,365,633, and WO 01/27060, and WO 01/162341; (xxiv) GLP-1 (glucagon-like peptide 1) agonists; (xxv) Topiramate 20 (Topimax®); (xxvi) phytopharm compound 57 (CP 644,673); (xxvii) ACC2 (acetyl-CoA carboxylase-2) inhibitors; (xxviii) β 3 (beta adrenergic receptor 3) agonists, such as AD9677/TAK677 (Dainippon/ Takeda), CL-316,243, SB 418790, BRL-37344, L-796568, BMS-196085, BRL-35135A, CGP12177A, BTA-243, GW 25 427353, Trecadrine, Zeneca D7114, and SR 59119A, and those disclosed in US Patent Application Nos. 5,705,515, US 5,451,677; and WO 01/74782, and WO 02/32897; (xxix) DGAT1 (diacylglycerol acyltransferase 1) inhibitors; (xxx) DGAT2 (diacylglycerol acyltransferase 2) inhibitors; (xxxi) FAS (fatty acid synthase)

inhibitors, such as Cerulenin and C75; (xxxii) PDE (phosphodiesterase) inhibitors, such as theophylline, pentoxifylline, zaprinast, sildenafil, amrinone, milrinone, cilostamide, rolipram, and cilomilast; (xxxii) thyroid hormone
5 β agonists, such as KB-2611 (KaroBioBMS), and those disclosed in WO 02/15845; and Japanese Patent Application No. JP 2000256190; (xxxiii) UCP-1 (uncoupling protein 1), 2, or 3 activators, such as phytanic acid, 4-[(E)-2-(5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthale
10 nyl)-1-propenyl]benzoic acid (TTNPB), and retinoic acid; and those disclosed in WO 99/00123; (xxxiv) acyl-estrogens, such as oleoyl-estrone, disclosed in del Mar-Grasa, M. et al., Obesity Research, 9:202-9 (2001); (xxxv) glucocorticoid antagonists; (xxxvi) 11β HSD-1 (11-beta hydroxy steroid
15 dehydrogenase type 1) inhibitors, such as BVT 3498, BVT 2733, and those compounds disclosed in WO 01/90091, WO 01/90090, WO 01/90092; (xxxvii) SCD-1 (stearoyl-CoA desaturase-1) inhibitors; (xxxviii) dipeptidyl peptidase IV (DP-IV) inhibitors, such as isoleucine thiazolidide, valine
20 pyrrolidide, NVP-DPP728, LAF237, P93/01, TSL 225, TMC-2A/2B/2C, FE 999011, P9310/K364, VIP 0177, SDZ 274-444; and the compounds disclosed in WO 03/004498, WO 03/004496, EP 1 258 476, WO 02/083128, WO 02/062764, WO 03/000250, WO 03/002530, WO 03/002531, WO 03/002553, WO 03/002593, WO 03/000180, and WO
25 03/000181; (xxxviii) lipase inhibitors, such as tetrahydrolipstatin (orlistat/Xenical®), Triton WR1339, RHC80267, lipstatin, teasaponin, and diethylumbelliferyl phosphate, FL-386, WAY-121898, Bay-N-3176, valilactone, esteracin, ebelactone A, ebelactone B, and RHC 80267, and those

disclosed in WO 01/77094, and U.S. Patent Nos. 4,598,089.
4,452,813, 5,512,565, 5,391,571, 5,602,151, 4,405,644,
4,189,438, and 4,242,453; (xxxix) fatty acid transporter
inhibitors; (xxxx) dicarboxylate transporter inhibitors;
5 (xxxxi) glucose transporter inhibitors; (xxxxii) phosphate
transporter inhibitors; (xxxxiii) melanocortin agonists, such
as Melanotan II or those described in WO 99/64002 and WO
00/746799; (xxxxiv) melanin concentrating hormone
antagonists; (xxxxv) galanin antagonists; (xxxxvi) CCK
10 agonists; (xxxxvii) corticotropin-releasing hormone agonists;
and (xxxxviii) phosphodiesterase-3B (PDE3B) inhibitors; and
the like.

The above combinations include combinations of a
composition of the present invention not only with one other
15 active compound, but also with two or more other active
compounds. Non-limiting examples include combinations of the
compositions of the present invention with one, two or more
active compounds selected from lipid-lowering agents, and
anti-hypertensive agents. Combinations of the compositions of
20 the present invention with one, two or more active compounds
selected from lipid lowering agents, and anti-diabetic agents
are useful to treat, control or prevent metabolic syndrome. In
particular, compositions comprising an anti-obesity agent, an
anti-hypertensive agent, in addition to an anti-diabetic agent
25 and/or a lipid lowering agent will be useful to synergistically
treat, control or prevent metabolic syndrome.

EXAMPLES

The following Examples and Reference Examples are provided to illustrate the present invention more concretely, but they should not be construed as limiting the invention in any way.

5 The mass spectrum was determined by electron spray ionization (ESI) method using QuattroII (Product of Micromass Ltd.).

Unless otherwise noted, melting point was disclosed in this specification without correction.

10

Example 1

Preparation of 5,6-dichloro-2-[1-methylsulfonyl-spiro[indoline-3,4'-piperidin]-1'-yl]benzimidazole

(1) Preparation of 2,5,6,-trichloro-N-tetrahydropyranyl-
15 benzimidazole

To a solution of 2,5,6-trichlorobenzimidazole (2.07 g, 9.33 mmol) in tetrahydrofuran (20 mL) were added 3,4-dihydro-2H-pyran (2.55 mL, 28.0 mmol) and camphor sulfonic acid (216 mg, 0.93 mmol). The mixture was stirred at room
20 temperature for 3 hours, and the reaction mixture was neutralized by addition of 0.1M aqueous sodium hydroxide, then extracted with ethyl acetate. The organic layer was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by silica gel
25 column chromatography (C-300, hexane:ethyl acetate = 4 : 1) to give the title compound (487 mg, 9 %).

(2) Preparation of 5,6-dichloro-2-[1-methylsulfonyl-spiro[indoline-3,4'-piperidin]-1'-yl]benzimidazole

To a solution of 2,5,6-trichloro-N-tetrahydropyranyl-benzimidazole (213 mg, 0.70 mmol) and 1-methylsulfonyl-spiro[indoline-3,4'-piperidine] hydrochloride (168 mg, 0.56 mmol) in dioxane (10 mL) was added cesium carbonate (562 mg, 5 0.47 mmol). The mixture was stirred at 150°C for 14 hours in a sealed tube and then cooled. After addition of water, the reaction mixture was extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and concentrated in vacuo. The resulting residue was dissolved in methanol (10 mL), and 10 camphor sulfonic acid (100 mg, 0.70 mmol) was added thereto, then stirred at room temperature for 12 hours. The reaction mixture was concentrated in vacuo, and the residue was neutralized with 0.1M aqueous sodium hydroxide, then extracted with chloroform. The organic layer was washed with brine, dried 15 over anhydrous sodium sulfate, and concentrated in vacuo. The precipitate formed upon addition of ethyl acetate to the residue was collected by filtration and dried to give the title compound (86 mg, 34 %).

¹HNMR (300 MHz, DMSO-d₆, δ ppm): 1.73-1.79 (2H, m), 1.85-1.96 20 (2H, m), 3.05 (3H, s), 3.18-3.30 (2H, m), 3.96 (2H, s), 4.08-4.18 (2H, m), 7.00-7.05 (1H, m), 7.20-7.32 (5H, m); mass spectrum (ESI): 452 (M+H)

The following compound was prepared according to the 25 procedure described in Example 1.

Example 2

2-[1-Methylsulfonylspiro[indoline-3,4'-piperidin]-1'-yl]-5-chlorobenzimidazole

¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.72-1.80 (2H, m), 1.87-1.98

(2H, m), 3.06 (3H, s), 3.12-3.28 (2H, m), 3.96 (2H, s), 4.08-4.17 (2H, m), 6.85-7.30 (7H, m), 11.51 (1H, s); mass spectrum(ESI):418(M+H)

5 Example 3

8-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-1'-yl]-2-phenylpurine

¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.87(2H, d, J=13.8Hz), 2.15 (2H, m), 3.62 (4H, m), 4.20 (2H, brd, J=12.6Hz), 7.24 (3H, 10 m), 7.43 (4H, m), 8.31 (1H, t, J=3.9Hz), 8.36 (2H, dd, J=6.9Hz), 8.50 (1H, brs); mass spectrum (ESI) : 411(M+H)

Example 4

Preparation of trans-5-methoxycarbonyl-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]benzimidazole

(1) Preparation of trans-N-[2-amino-5-methoxycarbonyl-phenyl]-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxamide

20 To a solution of trans-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxylic acid (20 mg, 0.081 mmol) and methyl 3,4-diaminobenzoate (20 mg, 0.12 mmol) in pyridine (1 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (25 mg, 0.13 mmol). The 25 mixture was stirred at room temperature for one hour under a nitrogen atmosphere, and then concentrated. After addition of saturated aqueous sodium bicarbonate to the residue, the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and concentrated

in vacuo. The residue was washed with ethyl acetate-hexane, and dried to give the title compound (30.4 mg, 95 %).

(2) Preparation of trans-5-methoxycarbonyl-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]benzimidazole

To a solution of trans-N-[2-amino-5-methoxycarbonyl-phenyl]-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxamide (27.7 mg, 0.070 mmol) in xylene (2 mL) was added a catalytic amount of p-toluenesulfonic acid monohydrate. The mixture was heated under reflux for 3 hours, cooled to room temperature, and concentrated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform : methanol=20 : 1) to give the title compound (20 mg, 75 %).

¹HNMR (300 MHz, CD₃OD, δ ppm) : 1.78-2.55 (8H, m), 3.25-3.44 (1H, m), 3.93 (3H, s), 7.51-7.58 (4H, m), 7.85 (1H, dd, J=1.0Hz, 7.6Hz), 7.93 (1H, dd, J=1.6Hz, 8.6Hz), 8.27 (1H, s); mass spectrum (ESI) : 377(M+H)

The following compound was prepared according to the procedure described in Example 4.

Example 5

Trans-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-5-phenylbenzimidazole

¹HNMR (400 MHz, CDCl₃, δ ppm) : 1.80-2.02 (2H, m), 2.16-2.68 (6H, m), 3.30-3.44 (1H, m), 7.20-7.90 (9H, m), 8.85 (1H, d, J=4.8Hz), 8.97 (1H, s); mass spectrum (ESI) : 396(M+H)

Example 6

Preparation of trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-phenylpurine

(1) Preparation of trans-N-[4-amino-2-phenyl-5-pyrimidyl]-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxamide

To a solution of trans-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxylic acid (190 mg, 0.77 mmol) and 2-phenyl-4,5-diaminopyrimidine (130 mg, 0.70 mmol) in pyridine (3.5 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (25 mg, 0.13 mmol) at 0°C. The mixture was stirred at room temperature for 10 hours under a nitrogen atmosphere and then concentrated. After addition of saturated aqueous sodium bicarbonate to the residue, the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate and concentrated. The residue was purified by silica gel column chromatography (chloroform : methanol=10 : 1) to give the title compound (212 mg, 73 %).

20

(2) Preparation of trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-phenylpurine

To a solution of trans-N-[4-amino-2-phenyl-5-pyrimidyl]-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxamide (100 mg, 0.24 mmol) in pyridine (3.5 mL) was added phosphorus oxychloride (0.027 mL, 0.28 mmol). The mixture was stirred at 50°C for 12 hours under a nitrogen atmosphere. The reaction mixture was cooled to room temperature, and then concentrated in vacuo. The residue was

diluted with ethyl acetate, and the mixture was washed successively with saturated aqueous sodium bicarbonate and saturated brine, dried over anhydrous magnesium sulfate, and concentrated in vacuo. The resulting residue was purified by preparative thin-layer chromatography (chloroform:methanol= 10 : 1) to give the title compound (87 mg, 91 %).

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.70-1.90 (2H, m), 2.20-2.50 (6H, m), 3.25-3.40 (1H, m), 7.40-7.70 (6H, m), 7.89 (1H, d, J=7.8Hz), 8.40-8.50 (2H, m), 9.21 (1H, s); mass spectrum (ESI) : 397 (M+H)

The following compounds were prepared according to the procedure described in Example 6.

Example 7

15 Cis-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-phenylpurine

¹HNMR (300 MHz, CDCl₃, δ ppm): 1.80-2.00 (2H, m), 2.05-2.45 (6H, m), 2.95-3.15 (1H, m), 7.36 (1H, d, J=7.8Hz), 7.45-7.65 (4H, m), 7.65-7.80 (1H, m), 7.99 (1H, d, J=7.5Hz), 8.42-8.55 (2H, m), 9.16 (1H, s); mass spectrum (ESI) : 397 (M+H)

Example 8

Trans-2-(2-fluorophenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

25 ¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.60-1.90 (2H, m), 2.05-2.35 (6H, m), 3.00-3.10 (1H, m), 7.20-7.75 (6H, m), 7.88 (1H, d, J=7.5Hz), 8.09 (1H, t, J=6.0Hz), 9.28 (1H, s); mass spectrum (ESI) : 415 (M+H)

Example 9

Trans-2-[2-(difluoromethoxy)phenyl]-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.40-1.90 (2H, m), 2.00-2.55
5 (6H, m), 2.85-3.00 (1H, m), 6.70 (1H, t, J=7.5Hz), 7.20-7.75
(5H, m), 7.88 (1H, d, J=7.5Hz), 7.97 (1H, d, J=7.5Hz), 8.55-8.70
(1H, m), 9.25 (1H, s); mass spectrum (ESI) : 463(M+H)

Example 10

10 Trans-2-(2,6-difluorophenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.75-1.88 (2H, m), 2.21-3.03
(6H, m), 3.41-3.52 (1H, br), 7.25 (2H, t, J=7.8Hz), 7.50-7.80
(5H, m), 7.82 (1H, d, J=7.8Hz), 9.18 (1H, s); mass spectrum (ESI)
15 : 433(M+H)

Example 11

Trans-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-2-phenylpurine

20 ¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.50-1.90 (2H, m), 2.15-2.40
(4H, m), 3.10-3.24 (1H, m), 7.47-7.65 (3H, m), 7.77 (1H, d,
J=4.8Hz), 8.35-8.50 (2H, m), 8.85 (1H, d, J=5.1Hz), 8.95 (1H,
s), 9.23 (1H, s) ; mass spectrum (ESI) : 398 (M+H)

25 Example 12

Trans-2-(3-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine

¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.88-2.56 (8H, m), 3.38-3.50
(1H, m), 7.28-7.40 (1H, m), 7.50-7.65 (1H, m), 7.87 (1H, d,

J=4.9Hz), 8.16 (1H, dd, J=1.1Hz, 10.8Hz), 8.29 (1H, d, J=7.8Hz), 8.87 (1H, d, J=4.9Hz), 9.15 (1H, s), 9.19 (1H, s); mass spectrum (ESI) : 416 (M+H)

5 Example 13

Trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine

(1) Preparation of trans-N-[4-amino-2-(4-fluorophenyl)-5-pyrimidinyl]-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide

To a solution of trans-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxylic acid (4.96 g, 20.0 mmol) and 2-(4-fluorophenyl)-4,5-diaminopyrimidine (3.7 g, 18.1 mmol) in pyridine (60 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (4.5 g, 23.5 mmol) at 0°C. The mixture was stirred at room temperature for 4 hours under a nitrogen atmosphere, and water (500 mL) and ethyl acetate (120 mL) were added thereto. The resulting precipitate was collected by filtration and dried to give the title compound as a colorless solid (3.97 g, 51 %). The filtrate was extracted with ethyl acetate, and the extract was dried over anhydrous magnesium sulfate, then concentrated. The residue was suspended in ethyl acetate, stirred for a while, filtered off, and dried to further give the title compound (3.13 g, 39 %).

¹HNMR (400 MHz, DMSO-d₆, δ ppm) : 1.85-1.95 (2H, m), 2.00-2.30 (6H, m), 2.75-2.85 (1H, m), 6.82 (2H, brs), 7.25 (2H, t, J=7.2Hz), 7.84 (1H, d, J=5.2Hz), 8.25-8.35 (2H, m), 8.40 (1H, s), 8.86 (1H, d, J=5.2Hz), 9.11 (1H, s), 9.23 (1H, s); mass

spectrum (ESI) : 434 (M+H)

(2) Preparation of trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine

To a solution of trans-N-[4-amino-2-(4-fluorophenyl)-5-pyrimidinyl]-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxamide (6.89 g, 15.9 mmol) in pyridine (100 mL) was added phosphorus oxychloride (2.3 mL, 24.6 mmol) at 0°C. After stirring at 50°C for 16 hours under a nitrogen atmosphere, the reaction mixture was cooled down to room temperature and concentrated in vacuo. Ice-water was added to the residue, and the mixture was stirred for a while, then extracted with chloroform. The organic layer was dried over anhydrous magnesium sulfate, and concentrated in vacuo. The residue was purified by silica gel column chromatography (C-300, chloroform : tetrahydrofuran = 4 : 1) to give the title compound (hereinafter referred to as Compound No. 13) as a colorless solid (3.0 g, 45 %).

¹HNMR (400 MHz, CDCl₃, δ ppm) : 1.44-2.58 (8H, m), 3.30-3.50 (1H, m), 7.19 (2H, t, J=8.8Hz), 7.80 (1H, d, J=4.8Hz), 8.46 (2H, dd, J=5.6Hz, 8.8Hz), 8.88 (1H, d, J=4.8Hz), 8.96 (1H, s), 9.18 (1H, s); mass spectrum (ESI) : 416 (M+H)

(3) Preparation of a crystal (Form-A) of Compound No. 13

Tetrahydrofuran (1.5 L) was added to the compound (5.0 g) obtained in the above procedure (2). The mixture was dissolved by heating under reflux, and concentrated until the volume of the solvent was reduced to about 30 mL. The solution was cooled

to room temperature, and the precipitated crystal formed upon stirring was collected by filtration to give a colorless crystal (Form-A, 4.1 g) of Compound No. 13.

M.p. 285°C-288°C (Yanagimoto Seisakusho, MP-S3).

Powder X-ray diffraction

2θ	Intensity(cps)
8.0	242
11.1	342
11.5	273
12.6	397
14.9	1497
16.0	808
16.2	728
17.7	1435
18.2	430
18.5	208
19.9	482
22.3	1103
22.6	1187
22.9	802
23.6	365
25.2	837
25.7	223
27.5	1585
27.9	178
28.2	202
28.7	362
29.6	165
30.2	538
33.8	163
34.1	185
35.5	153
36.6	135
41.5	147
41.6	153
45.3	178

5

The above powder X-ray diffraction data were measured using an automated X-ray apparatus RINT-Ultima+system (2kW) (Rigaku International Corporation). The analysis was carried out as follows:

10

X-ray source : Cu

Tube voltage/tube current : 40 kV/30 mA

Monochromator : automated monochromator

Goniometer : wide angle goniometer

5 Scan step : 0.02 deg.

Scan velocity : 2.00 deg./min.

Divergence slit : 1 deg.

Scattering slit : 1 deg.

Receiving slit : 0.15 mm

10 Determination temperature : 5°C to 40°C

(4) Preparation of a crystal (Form-B) of Compound No. 13

The crystal (Form-A, 4.6 g) obtained in the above procedure (3) was added to tetrahydrofuran (1.5 L), and the mixture was dissolved by heating under reflux. The solution was then cooled to 0°C, and a solution of maleic acid (1.33 g) in ethanol (10 ml) was added thereto. After the solvent was concentrated in vacuo, the residue was washed with ethyl acetate to give a maleate (6.0 g) of Compound No. 13.

20 The maleate (6.0 g) was suspended in water (300 ml), and the suspension was vigorously stirred at room temperature for 9 hours. The resulting precipitate was collected by filtration to give a colorless crystal (Form-B, 4.4 g) of Compound No. 13. M.p. 246°C-252°C (Yanagimoto Seisakusho, MP-S3).

Powder X-ray diffraction

2 θ	Intensity(cps)
9.5	447
10.8	1197
13.8	243
14.1	438
14.4	413
15.0	1408
16.0	457
16.1	528
16.9	795
17.2	1550
17.7	285
19.4	332
19.7	403
20.2	755
22.4	367
22.6	473
23.1	613
23.2	520
24.9	245
26.1	2600
26.5	695
27.1	527
27.8	657
28.3	488
28.8	457
29.8	438
30.3	350
31.7	250
32.0	333
48.8	277

The above X-ray diffraction data were measured under the same conditions as in Example 13(3) except that the tube current
5 was changed to 40 mA.

(5) Preparation of a crystal (Form-C) of Compound No. 13

The crystal (Form-A, 680 mg) of Compound No. 13 obtained in the above procedure of (3) was suspended in water (35 mL).
10 The suspension was stirred while heated under reflux for 15

hours, and the resulting precipitate was collected by filtration to give a colorless crystal (Form-C, 650 mg) of Compound No. 13.

M.p. 267°C-270°C (Mettler hot stage FP82).

5

Powder X-ray diffraction

2 θ	Intensity(cps)
9.7	652
11.6	328
12.6	375
13.3	907
14.2	638
15.5	692
15.9	787
18.3	3808
18.7	1475
18.9	798
19.5	337
19.8	927
20.2	1453
21.2	705
23.9	733
24.7	4958
25.3	400
25.6	1140
25.9	538
26.6	390
27.1	668
27.3	375
28.3	2590
28.7	2395
29.9	805
31.4	927
32.2	363
34.0	393
41.1	378
43.7	330

The above X-ray diffraction data were measured under the same conditions as in Example 13(3).

10 (6) Preparation of a crystal (Form-C) of Compound No. 13

(Alternative method of the procedure (5))

- (6-1) 2-Propanol (20 mL) and 1N aqueous potassium hydroxide (2.6 mL) were successively added to Compound No. 13 (1.0 g) at 0°C, followed by stirring at the same temperature for 30 minutes.
- 5 After dropwise addition of 0.15N hydrochloric acid (20 mL) to the reaction solution, the resulting precipitate was collected by filtration, and then dried to give a colorless solid (740 mg).
- 10 (6-2) The solid (380 mg) obtained above was suspended in 1N hydrochloric acid (10 mL), and the suspension was stirred at room temperature for 12 hours. The resulting precipitate was collected by filtration to give a colorless crystal (Form-C, 290 mg) of Compound No. 13.

15

The following compounds were prepared according to the procedure described in Example 6.

Example 14

20 Trans-2-(2-fluorophenyl)-8-[3-oxospiro[4-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-yl]purine

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.60-1.90 (2H, m), 2.10-2.50 (6H, m), 2.97-3.10 (1H, m), 7.20-7.45 (6H, m), 7.45-7.63 (2H, m), 7.93 (1H, d, J=7.8Hz), 8.12 (1H, t, J=7.8Hz), 8.92 (1H, d, J=4.5Hz), 9.28 (1H, s); mass spectrum (ESI) : 416 (M+H)

25

Example 15

Trans-2-(2,4-difluorophenyl)-8-[3-oxospiro[4-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-yl]purine

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.58-2.60 (8H, m), 3.10-3.80

(1H, m), 6.84-7.14 (2H, m), 7.78 (1H, d, J=4.8Hz), 8.17 (1H, dd, J=8.4Hz, J=14.8Hz), 8.87 (1H, d, J=4.8Hz), 8.94 (1H, s), 9.20 (1H, s); mass spectrum (ESI) : 434 (M+H)

5 Example 16

Trans-2-(4-fluorophenyl)-8-[3-oxospiro[5-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine

¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.80-2.60 (8H, m), 2.96-3.48 (1H, m), 7.22-7.42 (2H, m), 7.81 (1H, d, J=4.2Hz), 8.38-8.54
10 (2H, m), 8.84 (1H, d, J=4.2Hz), 9.07(2H, s), 9.10 (2H, s); mass spectrum (ESI) : 416 (M+H)

Example 17

Preparation of Trans-2-(2,5-difluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine

(1) Working up in the same manner as Example 13 by use of trans-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexane]-4'-carboxylic acid (24.0 g, 97.1 mmol) and 2-(2,5-
20 difluorophenyl)-4,5-diaminopyrimidine (19.4 g, 87 mmol), there was obtained the title compound (hereinafter referred to as Compound No. 17) as a colorless solid (11.3 g, 30 %).

¹HNMR(300 MHz, DMSO-d₆, δ ppm) : 1.83-2.02 (8H, m), 3.38-3.49 (1H, br), 7.31-7.48 (2H, m), 7.78-7.92 (2H, m), 8.88 (1H, d, J=4.9Hz), 9.19 (2H, s); mass spectrum (ESI) : 434 (M+H)
25

(2) Preparation of a crystal (Form-A) of Compound No. 17

Ethanol (390 mL) was added to Compound No. 17 (3.9 g) obtained in the above procedure (1). The mixture was dissolved

by heating under reflux, and concentrated until the volume of the solvent was reduced to 120 mL. The solution was cooled down to room temperature, and stirred for 12 hours. The resulting precipitate was collected by filtration to give a colorless
5 crystal (Form-A, 3.6 g) of Compound No.17.
M.p. 265°C-270°C (Mettler hot stage FP82).

Powder X-ray diffraction

2 θ	Intensity(cps)
9.1	387
14.3	448
15.2	155
16.6	1155
17.2	263
18.0	1147
18.3	315
18.8	933
19.1	1002
19.3	1582
19.7	628
21.0	287
21.8	583
22.3	303
22.9	997
23.7	407
24.2	428
24.6	518
24.8	802
25.3	368
26.0	270
27.7	278
28.4	365
29.1	245
29.6	243
31.0	393
33.3	235
33.7	175
36.7	192
38.2	162

The above powder X-ray diffraction data were measured in
10 the same manner as Example 13(3).

(3) Preparation of a crystal (Form-B) of Compound No. 17

- 2-Propanol (10 mL) and 1N aqueous potassium hydroxide (1.2 mL) were successively added to Compound No. 17 (500 mg) at 0°C, followed by stirring for 30 minutes. After dropwise addition of 0.15N hydrochloric acid (10 mL), the mixture was further stirred for 30 minutes. The resulting precipitate was collected by filtration to give a colorless crystal (Form-B, 390 mg) of Compound No. 17.
- 10 M.p. 223°C-227°C (Mettler hot stage FP82).

Powder X-ray diffraction

2 θ	Intensity(cps)
10.2	103
11.5	292
12.6	708
13.3	105
14.8	202
15.5	153
16.4	137
16.9	110
18.2	212
18.9	92
19.0	122
19.3	82
19.4	80
20.1	247
20.2	213
21.5	113
23.3	282
25.3	57
25.9	368
26.3	75
26.9	102
27.2	203
27.6	338
27.9	83
28.1	68
30.4	70
31.6	55
32.0	70
33.2	58

The above powder X-ray diffraction data were measured under the same conditions as Example 13(3).

5 The following compounds were prepared according to the procedure described in Example 6.

Example 18

Trans-2-(5-chloro-2-fluorophenyl)-8-[3-oxospiro[6-aza-
isobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine

10 ¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.70-1.95 (2H, m), 2.20-2.60 (6H, m), 3.20-3.40 (1H, m), 7.10-7.30 (1H, m), 7.40-7.55 (1H, m), 7.78 (1H, d, J=5.1Hz), 8.10-8.25 (1H, m), 8.87 (1H, d, J=5.1Hz), 8.93 (1H, s), 9.27 (1H, s); mass spectrum (ESI) : 450 (M+H)

15

Example 19

Trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-
cyclohexan]-4'-yl]-2-phenylpurine

20 ¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.90-2.06 (2H, m), 2.10-2.28 (4H, m), 2.29-2.45 (2H, m), 3.35-3.45 (1H, m), 7.42-7.56 (3H, m), 7.70 (1H, d, J=3.3Hz), 8.44 (2H, dd, J=7.8Hz, J=1.8Hz), 8.82 (1H, s), 9.12 (1H, s); mass spectrum (ESI) : 416 (M+H)

Example 20

25 Trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-
cyclohexan]-4'-yl]-2-(2-fluorophenyl)purine

¹HNMR (400MHz, DMSO-d₆, δ ppm) : 1.90-2.70 (8H, m), 3.08-3.60 (1H, m), 7.24-7.42 (2H, m), 7.42-7.60 (1H, m), 7.69 (1H, s), 8.00-8.14 (1H, m), 8.80 (1H, s), 9.13 (1H, s); mass spectrum

(ESI) : 434 (M+H)

Example 21

Trans-2-(2,4-difluorophenyl)-8-[5-fluoro-3-oxo-

5 spiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine

¹HNMR (400 MHz, DMSO-d₆, δ ppm) : 1.90-2.06 (2H, m), 2.06-2.60 (6H, m), 3.16-3.50 (1H, m), 7.14-7.30 (1H, m), 7.30-7.42 (1H, m), 7.69 (1H, s), 8.02-8.18 (1H, m), 8.79 (1H, s), 9.12 (1H, s); mass spectrum (ESI) : 459 (M+H)

10

Example 22

Trans-2-chloro-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-

isobenzofuran]-4-yl]purine

¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.73-1.82 (2H, m), 1.96-2.20 (4H, m), 2.23-2.35 (2H, m), 3.14-3.32 (1H, m), 7.56-7.65 (1H, m), 7.76-7.87 (3H, m), 8.82 (1H, s); mass spectrum (ESI) : 355, 357 (M+H)

Example 23

20 Preparation of trans-2-(6-fluoropyridin-2-yl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

(1) Preparation of trans-2-chloro-8-[3'-

oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-7-

(2-trimethylsilylethoxymethyl)purine

25 To a solution of trans-2-chloro-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine (205 mg, 0.49 mmol) in dimethylformamide (5 mL) were added 2-(trimethylsilyl)ethoxymethyl chloride (0.12 mL, 0.59 mmol) and 60% sodium hydride (30 mg, 0.75 mmol). The mixture was

stirred at 50°C for 3 days, cooled to room temperature, and diluted with ethyl acetate and water. The organic layer was washed successively with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by silica gel column chromatography (C-300, hexane : ethyl acetate = 3 : 2) to give the title compound (115mg, 41%).

(2) Preparation of trans-2-(6-fluoropyridine-2-yl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

To a solution of trans-2-chloro-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-7-(2-trimethylsilylethoxymethyl)purine (100 mg, 0.21 mmol) in toluene (2 mL) were added 6-fluoro-2-trimethylstannylpyridine (80 mg, 0.21 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride (17 mg, 0.02 mmol). The mixture was stirred under reflux at 120°C for 15 hours under a nitrogen atmosphere. After the reaction solution was cooled to room temperature, it was diluted with ethyl acetate and water. The organic layer was washed successively with water and saturated brine, dried over anhydrous magnesium sulfate, and then concentrated in vacuo. The residue was purified by silica gel column chromatography (C-300, hexane : ethyl acetate=2 : 3) to give the title compound (67 mg, 59 %).

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.67-1.95 (2H, m), 2.15-2.36 (6H, m), 3.40-3.52 (1H, m), 7.03-7.11 (1H, m), 7.42-7.63 (3H, m), 7.87 (1H, d, J=7.5Hz), 8.04 (1H, q, J=7.7Hz), 8.56 (1H, d,

J=7.7Hz), 9.23 (1H, s), 12.4 (1H, brs); mass spectrum (ESI) :
416 (M+H)

The following compounds were prepared according to the
5 procedure described in Example 23.

Example 24

Trans-2-(2-furyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-
isobenzofuran]-4-yl]purine

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.73-1.85 (2H, m), 2.25-2.47(6H,
10 m), 3.40-3.48 (1H, m), 6.64-6.66 (1H, m), 7.35 (1H, d, J=3.5Hz),
7.43-7.55 (2H, m), 7.60-7.68 (2H, m), 7.89 (1H, d, J=7.6Hz),
9.16 (1H, s), 11.17 (1H, s); mass spectrum (ESI) : 387 (M+H)

Example 25

15 Trans-2-(indol-5-yl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.70-1.82 (2H, m), 2.05-2.35
(6H, m), 3.13-3.22 (1H, m), 6.66 (1H, d, J=2.5Hz), 7.26-7.32
(1H, m), 7.55 (1H, d, J=8.5Hz), 7.79 (1H, d, J=5.0Hz), 8.24 (1H,
20 d, J=8.5Hz), 8.69 (1H, s), 8.85 (1H, d, J=5.0Hz), 9.00 (1H, s),
9.17 (1H, s); mass spectrum (ESI) : 437 (M+H)

Example 26

Preparation of trans-6-(4-fluorophenyl)-2-[3'-
25 oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-
yl]imidazo[4,5-b]pyridine

(1)Preparation of 2-amino-5-(4-fluorophenyl)-3-nitropyridine

To a suspension of 2-amino-5-bromo-3-nitropyridine (2.2

g, 10.1 mmol) in dimethoxyethane (20 mL) were added
4-fluorophenylboronic acid (4.83 g, 13.1 mmol),
tetrakis(triphenylphosphine)palladium (580 mg, 0.50 mmol) and
2N aqueous sodium carbonate solution (10 mL). The mixture was
5 stirred at 80°C for 9 hours under a nitrogen atmosphere. After
the reaction mixture was cooled to room temperature, it was
diluted with ethyl acetate to give a precipitate, which was
collected by filtration, washed with ethyl acetate and then
dried to give the title compound (1.76 g, 76 %).

10

(2) Preparation of 2,3-diamino-5-(4-fluorophenyl)pyridine

2-Amino-5-(4-fluorophenyl)-3-nitropyridine (200 mg,
0.86 mmol) was suspended in a mixture of methanol (5 mL), ethyl
acetate (5 mL), tetrahydrofuran (5 mL) and triethylamine (2 mL).
15 After addition of 10% palladium carbon (100 mg), the mixture
was stirred at room temperature for 1.5 hours under a hydrogen
atmosphere. The reaction mixture was filtered, and the
filtrate was concentrated in vacuo. The residue was purified
by silica gel column chromatography (C-300, chloroform :
20 methanol=50 : 1) to give the title compound (76 mg, 44 %) as
a white solid.

(3) Preparation of trans-N-[2-amino-5-(4-fluorophenyl)-3-
pyridyl]-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-
25 4-carboxamide

To a solution of 2,3-diamino-5-(4-fluorophenyl)pyridine
(76 mg, 0.37 mmol) and trans-3'-oxospiro[cyclohexane-
1,1'(3'H)-isobenzofuran]-4-carboxylic acid (92 mg, 0.37 mmol)
in pyridine (1 mL) was added

1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (86 mg, 0.45 mmol). The mixture was stirred at room temperature for 28 hours under a nitrogen atmosphere, and concentrated in vacuo. After addition of saturated aqueous sodium bicarbonate to the residue, the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and concentrated in vacuo. The residue was further washed with ethyl acetate and then dried to give the title compound (132 mg, 82 %).

(4) Preparation of trans-6-(4-fluorophenyl)-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]imidazo[4,5-b]pyridine

To a solution of trans-N-[2-amino-5-(4-fluorophenyl)-3-pyridyl]-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxamide (130 mg, 0.30 mmol) in pyridine (4 mL) was added phosphorus oxychloride (0.15 mL, 1.5 mmol). The mixture was stirred at 50°C for 16 hours. After the reaction mixture was cooled to room temperature, it was concentrated in vacuo. Aqueous ammonia was added to the residue, and the mixture was extracted with chloroform. The organic layer was dried over anhydrous magnesium sulfate, and concentrated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform : methanol=10 : 1) to give the title compound (32 mg, 32 %).

¹HNMR (300 MHz, CDCl₃, δ ppm) : 1.75-2.09 (2H, m), 2.23-2.75 (6H, m), 3.63 (1H, brs), 7.00-7.69 (7H, m), 7.88 (1H, d, J=7.4Hz), 8.29 (1H, brs), 8.53 (1H, brs), 13.2-13.6 (1H, br); mass spectrum (ESI) : 414 (M+H)

Example 27

Trans-5-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-b]pyridine

- 5 (1) Working up in the same manner as Example 13 by use of trans-3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-carboxylic acid (13.1 g, 53 mmol) and 2,3-diamino-6-(2-fluorophenyl)pyridine (10.8 g, 53 mmol), there was obtained the title compound (hereinafter referred to
10 as Compound No. 27) as a colorless solid (7.4 g, 34 %).

¹HNMR (400 MHz, DMSO-d₆, δ ppm) : 1.70-2.80 (8H, m), 2.80-3.68 (1H, m), 7.20-7.50 (2H, m), 7.61 (1H, d, J=7.2Hz), 7.76-8.02 (2H, m), 7.85 (1H, d, J=4.8Hz), 8.08 (1H, d, J=8.4Hz), 8.86 (1H, d, J=4.8Hz), 9.21 (1H, s); mass spectrum (ESI) : 415(M+H)

15

- (2) Preparation of a crystal (Form-A) of Compound No. 27

Ethyl acetate (2.0 L) was added to Compound No. 27 (6.1 g) obtained in the above procedure (1). The mixture was dissolved by heating to reflux, and concentrated until the
20 volume of the solvent was reduced to 30 mL. The solution was cooled down to room temperature, and stirred vigorously. The resulting precipitate was collected by filtration to give a colorless crystal (Form-A, 5.7 g) of Compound No 27.

M.p. 242°C-244°C (BUCHI Melting Point B-545).

25

Powder X-ray diffraction

2 θ	Intensity(cps)
8.0	1092
12.1	390
12.3	328
13.9	1273
15.2	312
15.4	320
16.2	937
17.4	537
18.3	177
19.4	303
20.2	288
20.3	355
20.9	760
21.3	580
21.8	690
22.5	167
23.4	193
24.4	552
25.3	258
25.9	165
26.0	167
26.5	458
26.6	527
27.0	472
27.9	222
28.1	217
28.9	168
29.1	190
29.3	215
29.6	213

The above powder X-ray diffraction data were measured under the same conditions as in Example 13(3).

5 (3) Preparation of a crystal (Form-B) of Compound No. 27

The colorless crystal (Form-A, 20 mg) of Compound No. 27 obtained in the above procedure was suspended in water (2 ml). The suspension was stirred while heated under reflux for 12 hours, and the resulting precipitate was collected by
10 filtration to give a colorless crystal (Form-B, 19 mg) of

Compound No. 27.

M.p. 255°C-256°C (BUCHI Melting Point B-545).

Powder X-ray diffraction

2 θ	Intensity(cps)
8.2	255
9.7	767
11.5	1162
12.8	358
13.2	323
14.4	250
14.9	330
15.3	797
16.5	490
18.1	2052
19.5	1180
20.0	1012
21.0	1552
23.1	475
23.4	652
24.4	2197
25.1	357
25.3	262
25.9	770
27.2	553
27.6	810
28.0	773
28.2	920
28.6	250
29.0	525
30.4	318
32.9	243
34.9	252
37.0	255
44.5	247

5 The above powder X-ray diffraction data were measured in the same manner as Example 13(3).

 The following compounds were prepared according to the procedure described in Example 26.

10 Example 28

Trans-6-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-c]pyridine

¹HNMR (400 MHz, DMSO-d₆, δ ppm) : 1.70-2.70 (8H, m), 2.70-3.60 (1H, m), 7.20-7.48 (3H, m), 7.48-8.12 (2H, m), 7.85 (2H, d, J=4.8Hz), 8.86 (1H, d, J=4.8Hz), 9.00 (1H, s), 9.16 (1H, s);
5 mass spectrum (ESI) : 415 (M+H)

Example 29

10 Trans-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-5-phenylimidazo[4,5-b]pyrazine

¹HNMR (400 MHz, CDCl₃, δ ppm) : 1.70-2.70 (8H, m), 3.22-3.60 (1H, m), 7.20-8.20 (9H, m), 8.70-9.00 (1H, m); mass spectrum (ESI) : 397 (M+H)

15 Example 30

Trans-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-6-phenylimidazo[4,5-b]pyridazine

¹HNMR (400 MHz, DMSO-d₆, δ ppm) : 1.70-1.90 (2H, m), 2.00-2.80 (6H, m), 3.20-3.40 (1H, m), 7.40-8.00 (7H, m), 8.10-8.40 (3H, m);
20 mass spectrum (ESI) : 397 (M+H)

Example 31

Trans-2-(2-fluorophenoxy)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

25 ¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.73-1.87 (2H, m), 2.04-2.25 (4H, m), 2.29-2.41 (2H, m), 3.25-3.50 (1H, m), 7.24-7.45 (4H, m), 7.59 (1H, d, J=7.6Hz), 7.64-7.80 (2H, m), 7.83 (1H, d, J=7.6Hz), 8.85 (1H, s), 13.28 (1H, brs); mass spectrum (ESI) : 431 (M+H)

Example 32

Trans-2-(2,6-difluorophenoxy)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

- 5 ¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.72-1.84 (2H, m), 2.03-2.25 (4H, m), 2.31-2.41 (2H, m), 3.23-3.49 (1H, m), 7.26-7.44 (3H, m), 7.59 (1H, d, J=7.3Hz), 7.65-7.80 (2H, m), 7.83 (1H, d, J=7.3Hz), 8.87 (1H, s), 13.38 (1H, brs); mass spectrum (ESI) : 449 (M+H)

10

Example 33

Trans-2-morpholino-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

- 15 ¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.72-1.86 (2H, m), 2.05-2.23 (4H, m), 2.24-2.38 (2H, m), 3.21-3.42 (1H, m), 3.65-3.74 (8H, m), 7.57-7.68 (2H, m), 7.73 (1H, d, J=7.7Hz), 7.84 (1H, d, J=7.7Hz), 8.67 (1H, s), 12.72 (1H, brs); mass spectrum (ESI) : 406 (M+H)

20 Example 34

Trans-2-(N-methylanilino)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine

- 25 ¹HNMR (300 MHz, DMSO-d₆, δ ppm) : 1.71-1.87 (2H, m), 2.02-2.22 (4H, m), 2.23-2.38 (2H, m), 3.20-3.41 (1H, m), 3.51 (3H, s), 7.14-7.23 (1H, m), 7.33-7.44 (4H, m), 7.60 (1H, t, J=7.4Hz), 7.66 (1H, d, J=7.4Hz), 7.74 (1H, t, J=7.4Hz), 7.84 (1H, d, J=7.4Hz), 8.66 (1H, s), 12.81 (1H, brs); mass spectrum (ESI) : 426 (M+H)

Example 35

Trans-2-(4-fluorophenyl)-8-[7-hydroxy-3-oxospiro[6-azaisobenzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine

¹HNMR(400MHz, CD₃OD, δppm): 2.22-2.50(2H, m), 2.50-2.80(6H, m),
5 3.42-3.70(1H, m), 6.62(1H, d, J=6.8Hz), 7.12-7.30(2H, m),
7.51(1H, d, J=6.8Hz), 8.40-8.58(2H, m), 8.98(1H, s); mass
spectrum(ESI): 432(M+H)

Example 36

10 Trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1
(3H), 1'-cyclohexan]-4'-yl]purine 6-oxide

¹HNMR(400MHz, DMSO-d₆, δppm): 1.80-2.02(2H, m), 2.02-2.28(4H,
m), 2.28-2.62(2H, m), 3.02-3.60(1H, m), 7.32(2H, t, J=8.8Hz),
7.82(1H, d, J=6.8Hz), 8.32(2H, d, J=6.8Hz), 8.42-8.54(2H, m),
15 8.84(1H, s), 9.08(1H, s), 9.18(1H, s); mass
spectrum(ESI): 432(M+H)

Example 37

20 Trans-2-(4-fluoro-2-hydroxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine

¹HNMR(400MHz, CD₃OD, δppm): 1.94-2.10(2H, m), 2.22-2.58(6H, m),
3.42-3.58(1H, m), 6.62-6.78(2H, m), 7.82-7.90(1H, m),
8.50-8.62(1H, m), 8.83(1H, d, J=4.8Hz), 9.04(1H, s), 9.09(1H,
s); mass spectrum(ESI): 432(M+H)

25

Example 38

Trans-2-(4-fluorophenyl)-6-hydroxy-8-[3-oxospiro[6-azaisobenzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine

¹HNMR(400MHz, CD₃OD, δppm): 1.94-2.10(2H, m), 2.22-2.46(6H, m),

3.22-3.40(1H, m), 7.29(2H, t, J=8.4Hz), 7.85(1H, d, J=4.4Hz),
8.08-8.18(2H, m), 8.83(1H, d, J=4.4Hz), 9.18(1H, s); mass
spectrum(ESI):432(M+H)

5 Example 39

Trans-2-(4-hydroxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H), 1'-cyclohexan]-4'-yl]purine

¹HNMR(400MHz, CD₃OD, δppm):1.94-2.10(2H, m), 2.26-2.58(6H, m),
3.52-3.62(1H, m), 7.14(2H, d, J=8.8Hz), 7.88(1H, d, J=5.2Hz),
10 8.35(2H, d, J=8.8Hz), 8.85(1H, d, J=5.2Hz), 9.05(1H, s),
9.13(1H, s); mass spectrum(ESI):414(M+H)

Example 40

Trans-2-(4-fluoro-3-hydroxyphenyl)-8-[3-oxospiro[6-azaisobe
15 nzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine

¹HNMR(400MHz, DMSO-d₆, δppm):1.80-2.70(8H, m), 3.00-3.54(1H,
m), 7.23(1H, d, J=7.6Hz), 7.80-8.00(2H, m), 8.07(1H, d,
J=7.6Hz), 8.86(1H, d, J=5.2Hz), 9.06(1H, s), 9.16(1H, s); mass
spectrum(ESI):432(M+H)

20

Example 41

Cis-2-(4-fluorophenyl)-8-[4'-hydroxy-3-oxospiro[6-azaiso-be
nzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine

¹HNMR(400MHz, CD₃OD, δppm):1.94-3.42(8H, m), 7.10-7.26(2H, m),
25 7.80-7.98(1H, m), 8.20-8.40(2H, m), 8.80-8.96(1H, m), 9.01(1H,
s), 9.09(1H, s); mass spectrum(ESI):432(M+H)

Reference Example 1

Preparation of 2-phenyl-4,5-diaminopyrimidine

2-phenyl-4-chloro-5-nitro-6-aminopyrimidine (367 mg, 1.46 mmol) which was produced according to the method described in the Journal of Chemical Society (C), 1408-1412 (1969) was suspended in a mixture of methanol (3 mL), ethyl acetate (2 mL) and triethylamine (0.21 mL). After addition of 10% palladium-carbon (40 mg), the mixture was stirred at room temperature for 2 hours under a hydrogen atmosphere and then filtered. The filtrate was concentrated in vacuo, and purified by silica gel column chromatography (C-300, chloroform : methanol=50 : 1) to give the title compound (175 mg, 64 %) as a brown solid.

^1H NMR (300 MHz, CDCl_3 , δ ppm) : 7.35-7.47 (3H, m), 7.97 (1H, s), 8.20-8.27 (1H, m); mass spectrum (ESI) : 187 (M+H)

15 Reference Example 2

Preparation of 2-phenyl-4,5-diaminopyrimidine

To a solution of 2-phenyl-5-nitropyrimidine (50.1 g, 250 mmol), which was produced according to the method described in the Journal of American Chemical Society, 78, 1434-1437 (1956), in dimethyl sulfoxide (1 L) were added trimethylammonium hydrazine iodide (106 g) and potassium t-butoxide (61.5 g). The mixture was stirred at room temperature for 2.5 hours, and then added to 1N hydrochloric acid. The resulting precipitate was collected by filtration to give a compound and the compound was dissolved in methanol (700 mL). After addition of 10% palladium carbon (8 g) to the solution, the mixture was stirred at room temperature overnight under a hydrogen atmosphere. The reaction mixture was filtered and the filtrate was concentrated in vacuo to give the title compound (25.9 g, 86 %) as a brown

solid.

Reference Example 3

Preparation of 2-phenyl-4,5-diaminopyrimidine

5 Zinc chloride (136 mg), potassium t-butoxide (505 mg) and
O-methylhydroxylamine monohydrochloride (125 mg) were
successively added to a solution of 2-phenyl-5-nitropyrimidine
(201 mg, 1.00 mmol), which was prepared according to the method
described in the Journal of American Chemical Society, 78,
10 1434-1437 (1956), in dimethyl sulfoxide (5 mL). The mixture
was stirred at room temperature for 40 minutes, and saturated
aqueous ammonium chloride was added thereto, then the mixture
was extracted with ethyl acetate. The organic layer was washed
with 5% aqueous sodium bicarbonate, dried over aqueous sodium
15 sulfate, and concentrated in vacuo. The residue was dissolved
in methanol (8 mL), and 10% palladium carbon (50 mg) was added
thereto. The mixture was stirred at room temperature for one
hour under a hydrogen atmosphere, and then filtered. The
filtrate was concentrated in vacuo to give the title compound
20 (148 mg, 80 %) as a brown solid.

Reference Example 4

Preparation of 2,3-diamino-6-(2-fluorophenyl)pyridine

(1) Preparation of 6-amino-2-bromo-5-nitropyridine

25 To a solution of commercially available 2,6-dichloro-5-
nitropyridine (20.0 g, 95.3 mmol) in ethanol (300 mL) was added
aqueous ammonia (60 mL), and the mixture was stirred at room
temperature for 10 hours. The resulting precipitate was
filtered, washed with ethanol, and dried in vacuo to give

6-amino-2-chloro-5-nitropyridine (13.8 g, 83 %). The compound thus obtained was suspended in an acetic acid solution (130 mL) of 30% hydrogen bromide. The suspension was stirred at 100°C for 26 hours, cooled to room temperature, and concentrated in vacuo. The resulting residue was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate and saturated brine, dried over anhydrous magnesium sulfate, and concentrated in vacuo. The residue thus obtained was recrystallized from ethyl acetate to give the title compound (14.93 g, 86 %).

(2) Preparation of 2,3-diamino-6-(2-fluorophenyl)pyridine

To a solution of 6-amino-2-bromo-5-nitropyridine (2.0 g, 9.2 mmol) and 2-fluorophenylboronic acid (2.57 g, 18 mmol) in dimethoxyethane (60 mL) were added 2M aqueous sodium carbonate solution (18 mL) and tetrakis(triphenylphosphine) palladium(0) (106 mg, 0.092 mmol). The mixture was stirred at 105°C for 16 hours under a nitrogen atmosphere. After the reaction mixture was cooled to room temperature, it was extracted with ethyl acetate. The organic layer was washed with saturated brine, dried over anhydrous magnesium sulfate, and concentrated in vacuo. The residue was dissolved in a mixture of ethyl acetate (30 mL) and methanol (30 mL). To the solution was added 10% palladium carbon (200 mg), and the mixture was stirred at room temperature for 14 hours under a hydrogen atmosphere. After removal of the palladium catalyst by filtration, the filtrate was concentrated in vacuo. The residue was purified by silica gel column chromatography (chloroform : methanol=30 : 1) to give the title compound (1.87 g, 100 %) as a reddish brown oil.

¹HNMR (300 MHz, CD₃OD, δ ppm) : 6.89-6.96 (2H, m), 7.06-7.12 (1H, m), 7.15-7.20 (1H, m), 7.25-7.30 (1H, m), 7.65-7.70 (1H, m); mass spectrum (ESI) : 204 (M+H)

5 Reference Example 5

Preparation of 4,5-diamino-2-(4-fluorophenyl)pyrimidine

(1) Preparation of 2-(4-fluorophenyl)-4,6-dihydroxy-5-nitropyrimidine

4,6-dihydroxy-2-(4-fluorophenyl)pyrimidine (11 g, 53
10 mmol) described in European patent No. 136976A was gradually added to fuming nitric acid (100 mL) at 0°C. The reaction temperature was raised to room temperature, and the mixture was stirred for 15 hours. The reaction mixture was poured onto ice-water (150 mL), and then neutralized with sodium carbonate
15 at 0°C. The resulting precipitate was collected by filtration and dried to give the objective compound as a solid (9.2 g, 68 %).

(2) Preparation of 4,6-dichloro-2-(4-fluorophenyl)-5-nitropyrimidine
20

Phosphorus oxychloride (17 mL) and N,N-dimethylaniline (4.21 mL) were added to 2-(4-fluorophenyl)-4,6-dihydroxy-5-nitropyrimidine (8.2 g, 32.6 mmol). The mixture was heated under reflux for 3 hours, cooled down to room temperature, and
25 concentrated in vacuo to give a residue, to which was added ice-water. The resulting precipitate was collected by filtration, washed with water, and dried to give the objective compound as a solid (9.0 g, 96 %).

(3) Preparation of 4-amino-6-chloro-2-(4-fluorophenyl)-5-nitropyrimidine

28% aqueous ammonia (3.1 mL) was added to a solution of 4,6-dichloro-2-(4-fluorophenyl)-5-nitropyrimidine (5.0 g, 17.4 mmol) in tetrahydrofuran (50 mL) at -50°C until the starting material disappeared. After addition of 3N hydrochloric acid (5 mL) to the reaction solution at the same temperature, the reaction temperature was raised and then the mixture was concentrated in vacuo. Water was added to the residue, and the resulting suspension was filtered, then dried to give the objective compound as a solid (4.05 g, 87 %).

(4) Preparation of 4,5-diamino-2-(4-fluorophenyl)pyrimidine

4-amino-6-chloro-2-(4-fluorophenyl)-5-nitropyrimidine (15.3 g, 57 mmol) was dissolved in a mixture of methanol (200 mL) and ethyl acetate (100 mL), and after adding thereto 10% palladium on carbon (2.88 g) and triethylamine (12 mL), the mixture was stirred in a hydrogen stream for 18 hours. The reaction mixture was filtered and the filtrate was concentrated in vacuo to give a residue, to which was added water. The mixture was extracted with ethyl acetate, and the organic layer was dried over anhydrous magnesium sulfate, then concentrated in vacuo. The residue was purified by silica gel column chromatography (C-300, chloroform : methanol = 20 : 1) to give the title compound as a brown solid (7.16 g, 68 %).

¹HNMR (400 MHz, DMSO-d₆, δ ppm) : 4.83 (2H, s), 6.36 (1H, s), 7.10-7.23 (2H, m), 7.64 (1H, s), 8.11-8.22 (2H, m); mass spectrum (ESI) : 205 (M+H)

Reference Example 6

Preparation of 4,5-diamino-2-(2,5-difluoro- phenyl)-
pyrimidine

Working up in the same manner as Reference Example 5, there
5 was obtained the title compound (59.7 g) as a yellow solid from
3,5-dihydroxy-2-(2,5-difluorophenyl)pyrimidine (125 g).

Mass spectrum (ESI) : 223 (M+H)

Formulation Example 1

10 20.0 g of the compound of Example 6, 417 g of lactose,
80 g of crystalline cellulose and 80g of partial alpha-starch
were blended with a V-cone blender. To the mixture was added
3.0 g of magnesium stearate, and the whole was blended. The
blended powder was compressed into 3000 tablets by conventional
15 procedure so that each tablet has a weight of 150 mg and a
diameter of 7.0 mm.

The content per one tablet(150 mg)

	The compound of Example 1	5.0 mg
20	Lactose	104.25 mg
	Crystalline cellulose	20.0 mg
	Partial alpha-starch	20.0 mg
	Magnesium stearate	0.75 mg

25 Formulation Example 2

10.8 g of hydroxypropylcellulose 2910 and 2.1 g of
polyethylene glycol 6000 were dissolved in 172.5 g of purified
water. To the solution was dispersed 2.1 g of titanium dioxide
to provide a coating liquid. 2,500 tablets separately prepared

according to Formulation Example 1 were subjected to spray-coating with the coating liquid using HICOATER-MINI to provide film coated tablets with a weight of 155 mg.

5 The content per one tablet(155 mg)

The tablet prepared in Formulation Example 1	150 mg
Hydroxypropylcellulose 2910	3.6 mg
Polyethylene glycol 6000	0.7 mg
Titanium dioxide	0.7 mg

10

Industrial Applicability


Compounds of the present invention (I) exhibit NPY antagonistic effects, show excellent pharmacokinetics such as transport into brain or transport to cerebrospinal fluid, etc., and are very safe. Thus, the compound of the present invention (I) are useful as agents for the treatment of various diseases related to NPY, for example, cardiovascular disorders such as angina, acute or congestive heart failure, myocardial infarction, hypertension, nephropathy, electrolyte abnormality, vasospasm, arteriosclerosis, etc., central nervous system disorders such as bulimia, depression, anxiety, seizure, epilepsy, dementia, pain, alcoholism, drug withdrawal, circadian rhythm disorders, schizophrenia, memory impairment, sleep disorders, cognitive impairment, etc., metabolic diseases such as obesity, diabetes, hormone abnormality, hypercholesterolemia, hyperlipidemia, gout, fatty liver, etc., genital or reproductive disorders such as infertility, preterm labor, sexual dysfunction, etc., gastro-intestinal disorders, respiratory disorder,

15
20
25

inflammatory diseases or glaucoma, and the like, also for example, atherosclerosis, hypogonadism, hyperandrogenism, polycystic ovary syndrome, hirsutism, gastro-intestinal motility disorder, obesity-related gastro-esophageal
5 reflux, obesity hypoventilation (Pickwickian syndrome), sleep apnea, inflammation, systemic inflammation of the vasculature, osteoarthritis, insulin resistance, bronchoconstriction, alcohol preference, metabolic syndrome, Alzheimer's disease, cardiac hypertrophy, left
10 ventricular hypertrophy, hypertriglyceridemia, low HDL cholesterol, cardiovascular disorders such as coronary heart disease (CHD), cerebrovascular disease, stroke, peripheral vascular disease, sudden death, gallbladder diseases, cancer (breast, endometrial, colon), breathlessness, hyperuricemia,
15 impaired fertility, low back pain, or increased anesthetic risk, and the like.

CLAIMS

1. A compound of the formula (I),



5 (wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower
10 alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$, and at least one of A, B, C and D is said methine group;

Ar^1 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl,
15 hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower
20 alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;

25 n is 0 or 1;

Q^1 and Q^2 are independently a single bond, oxygen, carbonyl or $-N(R^3)-$;

R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be
5 intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;

10 R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;

T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower
15 alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group;

X is $-N(SO_2R^4)-$, $-N(COR^5)-$ or $-CO-$;

Y is $-C(R^6)(R^7)-$, $-O-$ or $-N(R^8)-$, provided that the compound (I) when E is nitrogen, n is 0, X is $-CO-$ and Y is $-O-$
20 is excluded), and a salt, ester or N-oxide derivative thereof.

2. The compound as claimed in claim 1, wherein n is 0, X is $-N(SO_2R^4)-$ or $-N(COR^5)-$ and Y is $-C(R^6)(R^7)-$, or X is $-CO-$ and Y is $-O-$ or $-N(R^8)-$.

25

3. The compound as claimed in claim 1, wherein T, U, V and W are independently methine optionally substituted by halogen, lower alkyl, hydroxy or lower alkoxy.


4. The compound as claimed in claim 1, wherein at least one of T, U, V and W is nitrogen.

5. The compound as claimed in claim 1, wherein one of T, U, V and W is nitrogen, and one of the remainder is methine optionally substituted by halogen, lower alkyl, hydroxy or lower alkoxy.

6. The compound as claimed in claim 1, wherein X is -CO- and Y is -O- or -NH-.

7. The compound as claimed in claim 1, wherein X is -CO- and Y is -O-.

8. The compound of the formula (I-a) as claimed in claim 1,



wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R¹)R² and -Q¹-Ar¹, and at least one of A, B, C and D is said methine group;

Ar¹ is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting

of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl,
 5 lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower
 10 alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;


Q^1 and Q^2 are independently a single bond, oxygen, carbonyl or $-N(R^3)-$;

R^1 and R^2 are independently hydrogen or lower alkyl, or
 15 R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

R^4 is hydrogen, lower alkyl, aralkyl or aryl.

20 9. The compound of the formula (I-b) as claimed in claim 1,



wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent
 25 selected from the group consisting of halogen, cyano, lower

alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$, and at least one of A, B, C and D is said methine group;

5 Ar^1 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower
10 alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower
15 alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

Q^1 and Q^2 are independently a single bond, oxygen, carbonyl or $-N(R^3)-$;

R^0 is hydrogen or hydroxy;

20 R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

 T, U, V and W are independently methine or nitrogen, said
25 methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group.


10. The compound as claimed in claim 9, wherein each of T, U, V and W is unsubstituted methine.

11. The compound as claimed in claim 9, wherein one of T, U, V and W is nitrogen.

12. The compound as claimed in claim 9, wherein one of T, U, V and W is nitrogen, and one of the remainder is methine having a substituent selected from the group consisting of halogen, lower alkyl, hydroxy and lower alkoxy.

13. The compound as claimed in claim 9, wherein one of T, U, V and W is nitrogen, and one of the remainder is methine having a substituent selected from the group consisting of fluorine and hydroxy.

14. The compound of the formula (I-c) as claimed in claim 1,



wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R¹)R² and -Q¹-Ar¹, and at least one of A, B, C and D is said methine group;

Ar¹ is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino and -Q²-Ar²;

Ar² is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;

Q¹ and Q² are independently a single bond, oxygen, carbonyl or -N(R³)-;

R¹ and R² are independently hydrogen or lower alkyl, or R¹ and R², taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino; and

R³ is hydrogen or lower alkyl.

20

15. The compound as claimed in any one of claims 1, 8, 9, 10, 11, 12, 13 and 14, wherein Ar¹ is phenyl which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino and -Q²-Ar².

16. The compound as claimed in any one of claims 1, 8, 9, 10, 11, 12, 13 and 14, wherein Ar¹ is heteroaryl which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxy carbonyl, lower alkanoylamino and -Q²-Ar².
17. The compound as claimed in claim 16, wherein the heteroaryl is pyridyl.
18. The compound as claimed in claim 1, which is selected from the group consisting of
- 2-[1-methylsulfonylspiro[indoline-3,4'-piperidin]-1'-yl]-5-chlorobenzimidazole,
- 5,6-dichloro-2-[1-methylsulfonylspiro[indoline-3,4'-piperidin]-1'-yl]benzimidazole,
- 5-chloro-2-[1-ethylsulfonylspiro[indoline-3,4'-piperidin]-1'-yl]benzimidazole,
- trans-5-chloro-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-6-(trifluoromethyl)benzimidazole,
- trans-5-(4-fluorophenyl)-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]benzimidazole,
- trans-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-5-phenylbenzimidazole,
- trans-5-(3-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]benzimidazole,
- trans-5-(4-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-

- 1(3H),1'-cyclohexan]-4'-yl]benzimidazole,
trans-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-
4-yl]-5-(2-pyridyl)imidazo[4,5-b]pyridine,
trans-2-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-
5 4-yl]-6-(2-pyridyl)imidazo[4,5-b]pyridine,
trans-5-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-b]pyridine,
trans-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclo-
hexan]-4'-yl]-6-phenylimidazo[4,5-c]pyridine,
10 trans-6-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-c]pyridine,
trans-2-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclo-
hexan]-4'-yl]-5-phenylimidazo[4,5-b]pyrazine,
trans-5-(4-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-
15 1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-b]pyrazine,
trans-6-(4-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]imidazo[4,5-c]pyridazine,
trans-2-(2-fluorophenyl)-8-[3'-oxospiro[cyclohexane-
1,1'(3'H)-isobenzofuran]-4-yl]purine,
20 trans-2-(3-fluorophenyl)-8-[3'-oxospiro[cyclohexane-
1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(4-fluorophenyl)-8-[3'-oxospiro[cyclohexane-
1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-methoxyphenyl)-8-[3'-oxospiro[cyclohexane-
25 1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(4-methoxyphenyl)-8-[3'-oxospiro[cyclohexane-
1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-methylphenyl)-8-[3'-oxospiro[cyclohexane-
1,1'(3'H)-isobenzofuran]-4-yl]purine,

- trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-(2-trifluoromethylphenyl)purine,
trans-2-(2-chloro-4-fluorophenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
5 trans-2-(2-hydroxymethylphenyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-furyl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-(2-thienyl)purine,
10 trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-(2-pyrrolyl)purine,
trans-2-(3-fluoropyridin-6-yl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
15 trans-2-(2-chloropyridin-6-yl)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2-fluorophenoxy)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
trans-2-(2,6-difluorophenoxy)-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]purine,
20 trans-8-[3'-oxospiro[cyclohexane-1,1'(3'H)-isobenzofuran]-4-yl]-2-(1-piperidyl)purine,
trans-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-2-phenylpurine,
25 trans-2-(2-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(3-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-

- 1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2-chlorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(3-chlorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
5 1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(4-chlorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2-chloro-4-fluorophenyl)-8-[3-oxospiro[6-azaisoben
zofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
10 trans-2-(4-methoxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2-methylphenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2-difluoromethoxyphenyl)-8-[3-oxospiro[6-azaiso-
15 benzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(3-difluoromethoxyphenyl)-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclo-
hexan]-4'-yl]-2-(2-trifluoromethylphenyl)purine,
20 trans-2-(2,4-difluorophenyl)-8-[3-oxospiro[6-azaisobenzo-
furan-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2,5-difluorophenyl)-8-[3-oxospiro[6-azaisobenzo-
furan-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2-bromo-4-fluorophenyl)-8-[3-oxospiro[6-azaiso-
25 benzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(4-chloro-2-fluorophenyl)-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-8-[3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclo-
hexan]-4'-yl]-2-(3-quinolyl)purine,

- trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-2-phenylpurine,
trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]-2-(2-fluorophenyl)purine,
5 trans-2-(2,4-difluorophenyl)-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(2,5-difluorophenyl)-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-8-[7-hydroxy-3-oxospiro[6-azaisobenzofuran-1(3H),1'-cyclohexan]-4'-yl]purine,
10 5,6-dichloro-2-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-1'-yl]benzimidazole,
2-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-1'-yl]-5-phenylbenzimidazole,
15 8-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-1'-yl]-2-phenylpurine,
8-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-1'-yl]-2-(2-fluorophenyl)purine,
8-[3,4-dihydro-3-oxospiro[isoquinoline-1(2H),4'-piperidin]-1'-yl]-2-(4-fluorophenyl)purine,
20 trans-8-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-isoquinolin]-4-yl]-2-phenylpurine,
trans-2-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-isoquinolin]-4-yl]-5-phenylbenzimidazole,
25 trans-8-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-isoquinolin]-4-yl]-2-(2-fluorophenyl)purine,
trans-8-[3',4'-dihydro-3'-oxospiro[cyclohexane-1,1'(2'H)-isoquinolin]-4-yl]-2-(4-fluorophenyl)purine,
trans-2-(4-fluorophenyl)-8-[7-hydroxy-3-oxospiro[6-azaiso-


- benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H), 1'-cyclohexan]-4'-yl]purine 6-oxide,
trans-2-(4-fluoro-2-hydroxyphenyl)-8-[3-oxospiro[6-azaiso-
5 benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine,
trans-2-(4-fluorophenyl)-6-hydroxy-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine,
trans-2-(4-hydroxyphenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H), 1'-cyclohexan]-4'-yl]purine,
10 trans-2-(4-fluoro-3-hydroxyphenyl)-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine, and
cis-2-(4-fluorophenyl)-8-[4'-hydroxy-3-oxospiro[6-azaiso-
benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine.
- 15 19. The compound of claim 1, which is
trans-8-[3'-oxospiro[cyclohexane-1, 1'(3'H)-isobenzofuran]-
4-yl]-2-phenylpurine.
- 20 20. The compound of claim 1, which is
trans-2-(2-fluorophenyl)-8-[3'-oxospiro[cyclohexane-
1, 1'(3'H)-isobenzofuran]-4-yl]purine.
- 25 21. The compound of claim 1, which is
trans-2-(4-fluorophenyl)-8-[3-oxospiro[6-azaisobenzofuran-
1(3H), 1'-cyclohexan]-4'-yl]purine.
22. The compound of claim 1, which is
trans-2-(2,5-difluorophenyl)-8-[3-oxospiro[6-azaiso-
benzofuran-1(3H), 1'-cyclohexan]-4'-yl]purine.

23. The compound of claim 1, which is
trans-8-[5-fluoro-3-oxospiro[6-azaisobenzofuran-1(3H), 1'-
cyclohexan]-4'-yl]-2-(2-fluorophenyl)purine.

5

24. The compound of claim 1, which is
trans-5-(2-fluorophenyl)-2-[3-oxospiro[6-azaisobenzofuran-
1(3H), 1'-cyclohexan]-4'-yl]imidazo[4, 5-b]pyridine.

10 25. A process for preparing a compound of the formula (I-1):



(wherein A, B, C and D are independently methine or nitrogen,
said methine being optionally substituted by a substituent
selected from the group consisting of halogen, cyano, lower
15 alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower
alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower
alkylsulfonyloxy, -N(R¹)R² and -Q¹-Ar¹, and at least one of A,
B, C and D is said methine group;


Ar¹ is aryl or heteroaryl, any of which is optionally
20 substituted by a substituent selected from the group consisting
of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl,
hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower
alkoxy, halo-lower alkoxy, lower alkylthio, lower
alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxy carbonyl,
25 lower alkanoylamino and -Q²-Ar²;

Ar² is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl; Q¹ and Q² are independently a single bond, oxygen, carbonyl or -N(R³)-;

R¹ and R² are independently hydrogen or lower alkyl, or R¹ and R², taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group; and

n, R³, X and Y have each the same meaning as defined below), or a salt, ester or N-oxide derivative thereof, which comprises reacting a compound of the formula (II):



(wherein a, b, c and d are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R^{1p})R^{2p}, -Q^{1p}-Ar^{1p} and optionally protected hydroxy, and at least one of a, b, c and d is said methine group;

Ar^{1p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, lower alkyl, halo-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino, -Q^{2p}-Ar^{2p}, optionally protected hydroxy, optionally protected hydroxy-lower alkyl and optionally protected carboxyl;

Ar^{2p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl, aryl, optionally protected hydroxy-lower alkyl and optionally protected hydroxy;

L¹ is a leaving group;


P is an imino-protecting group;

Q^{1p} and Q^{2p} are independently a single bond, oxygen, optionally protected carbonyl or -N(R³)-;

R^{1p} and R^{2p} are independently an amino-protecting group, an imino-protecting group, hydrogen or lower alkyl, or R^{1p} and R^{2p}, taken together, form lower alkylene which may be intervened by oxygen, sulfur or optionally protected imino; and

R³ is hydrogen or lower alkyl)

with a compound of the formula (III) or its salt:



(III)

(wherein n is 0 or 1;

R^4 is lower alkyl, aralkyl or aryl;


R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;

5 R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;

t, u, v and w are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower
10 alkyl, lower alkoxy, halo-lower alkoxy and optionally protected hydroxy, and at least two of t, u, v and w are said methine group;


X is $-\text{N}(\text{SO}_2\text{R}^4)-$, $-\text{N}(\text{COR}^5)-$ or $-\text{CO}-$;

Y is $-\text{C}(\text{R}^6)(\text{R}^7)-$, $-\text{O}-$ or $-\text{N}(\text{R}^8)-$, provided that the compound (III) when n is 0, X is $-\text{CO}-$ and Y is $-\text{O}-$ is excluded),
15 and optionally removing the protecting group(s) from the resulting compound of the formula (IV-1):



(wherein a, b, c, d, n, P, t, u, v, w, X and Y have each the same meaning as defined above) and/or oxidizing nitrogen
20 atom(s) of the compound(IV-1).

26. A process for preparing a compound of the formula (I-2):



(wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower
 5 alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R¹)R² and -Q¹-Ar¹, and at least one of A, B, C and D is said methine group;

Ar¹ is aryl or heteroaryl, any of which is optionally
 10 substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl,
 15 lower alkanoylamino and -Q²-Ar²;

Ar² is aryl or heteroaryl, any of which is optionally
 substituted by a substituent selected from the group consisting
 of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower
 alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower
 20 alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

Q¹ and Q² are independently a single bond, oxygen, carbonyl
 or -N(R³)-;


R⁰ is hydrogen or hydroxy;

R¹ and R² are independently hydrogen or lower alkyl, or
 25 R¹ and R², taken together, form lower alkylene which may be

intervened by oxygen, sulfur or imino;

T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group;

n, R³, X and Y have each the same meaning as defined below), or a salt, ester or N-oxide derivative thereof, which comprises reacting a compound of the formula (V):



10

(wherein a, b, c and d are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R^{1p})R^{2p}, -Q^{1p}-Ar^{1p} and optionally protected hydroxy, and at least one of a, b, c and d is said methine group;

Ar^{1p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, lower alkyl, halo-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino, -Q^{2p}-Ar^{2p}, optionally protected hydroxy, optionally protected hydroxy-lower alkyl and optionally protected carboxyl;


Ar^{2p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting

of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl, aryl, optionally protected hydroxy-lower alkyl and optionally protected hydroxy;

5 Q^{1p} and Q^{2p} are independently a single bond, oxygen, optionally protected carbonyl or $-N(R^3)-$;

R^{1p} and R^{2p} are independently an amino-protecting group, an imino-protecting group, hydrogen or lower alkyl, or R^{1p} and R^{2p} , taken together, form lower alkylene which may be intervened
10 by oxygen, sulfur or optionally protected imino; and

R^3 is hydrogen or lower alkyl) or its salt with a compound of the formula (VI):



(wherein n is 0 or 1;

15 R^{Op} is hydrogen or optionally protected hydroxy;

R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;


R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl,
20 aralkyl or aryl;

t, u, v and w are independently methine or nitrogen, the said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy and
25 optionally protected hydroxy, and at least two of t, u, v and

w are said methine group;


X is $-\text{N}(\text{SO}_2\text{R}^4)-$, $-\text{N}(\text{COR}^5)-$ or $-\text{CO}-$;

Y is $-\text{C}(\text{R}^6)(\text{R}^7)-$, $-\text{O}-$ or $-\text{N}(\text{R}^8)-$ or its salt, subjecting the resulting compound of the formula (VII):




5

(wherein a, b, c, d, n, R^{Op} , t, u, v, w, X and Y have each the same meaning as defined above) to intramolecular dehydrative ring closure to give a compound of the formula (VIII):



10 (wherein a, b, c, d, n, R^{Op} , t, u, v, w, X and Y have each the same meaning as defined above), and optionally removing the protecting group(s) from the compound (VIII) and/or oxidizing nitrogen atom(s) of the compound (VIII).

15 27. A process for preparing a compound of the formula (I-2):



(wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$, and at least one of A, B, C and D is said methine group;

Ar^1 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoylamino and aryl;

Q^1 and Q^2 are independently a single bond, oxygen, carbonyl or $-N(R^3)-$;

R^0 is hydrogen or hydroxy;

R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

T, U, V and W are independently methine or nitrogen, the said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy,


and at least two of T, U, V and W are said methine group;

n, R³, X and Y have each the same meaning as defined below),

or a salt, ester or N-oxide derivative thereof,

which comprises reacting a compound of the formula (V):

5



(wherein a, b, c and d are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, 10 -N(R^{1p})R^{2p}, -Q^{1p}-Ar^{1p} and optionally protected hydroxy, and at least one of a, b, c and d is said methine group;


Ar^{1p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, lower alkyl, halo-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino, -Q^{2p}-Ar^{2p}, optionally 15 protected hydroxy, optionally protected hydroxy-lower alkyl and optionally protected carboxyl;

Ar^{2p} is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl, aryl, optionally protected hydroxy-lower alkyl 20 and optionally protected hydroxy;

Q^{1p} and Q^{2p} are independently a single bond, oxygen, optionally protected carbonyl or $-N(R^3)-$;

R^{1p} and R^{2p} are independently an amino-protecting group, an imino-protecting group, hydrogen or lower alkyl, or R^{1p} and
 5 R^{2p} , taken together, form lower alkylene which may be intervened by oxygen, sulfur or optionally protected imino;

R^3 is hydrogen or lower alkyl) or its salt with a compound of the formula (IX):



10 (wherein n is 0 or 1;

R^{Op} is hydrogen or optionally protected hydroxy

R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;

15 R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;


t, u, v and w are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower
 20 alkyl, lower alkoxy, halo-lower alkoxy, and optionally protected hydroxy, and at least two of t, u, v and w are said methine group;

X is $-N(SO_2R^4)-$, $-N(COR^5)-$ or $-CO-$;

Y is $-C(R^6)(R^7)-$, $-O-$ or $-N(R^8)-$, or its salt,

25 and optionally removing the protecting group(s) from the

resulting compound of the formula (VIII):



lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower
5 alkyllamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;


Q^2 is a single bond, oxygen, carbonyl or $-N(R^3)-$;

R^1 and R^2 are independently hydrogen or lower alkyl, or
10 R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least
15 two of T, U, V and W are said methine;

n, R^3 , X and Y have each the same meaning as defined below), or a salt, ester or N-oxide derivative thereof,

which comprises reacting a compound of the formula (X):



20

(wherein a^0 , b^0 , c^0 and d^0 are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower
25 alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower

alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy,
 -N(R^{1p})R^{2p} and optionally protected hydroxy, and at least one
 of a⁰, b⁰, c⁰ and d⁰ is said methine group having halogen atom
 or trifluoromethanesulfonyloxy group; E^p is nitrogen, methine
 5 or methine substituted by optionally protected hydroxy;

n is 0 or 1;

P is an imino-protecting group;

R^{1p} and R^{2p} are independently an amino-protecting group,
 an imino-protecting group, hydrogen or lower alkyl, or R^{1p} and
 10 R^{2p}, taken together, form lower alkylene which may be
 intervened by oxygen, sulfur or optionally protected imino;

R⁴ is lower alkyl, aralkyl or aryl;

R⁵ and R⁸ are independently hydrogen, lower alkyl, aralkyl
 or aryl;

15 R⁶ and R⁷ are independently hydrogen, hydroxy, lower alkyl,
 aralkyl or aryl;

t, u, v and w are independently methine or nitrogen, said
 methine being optionally substituted by a substituent selected
 from the group consisting of halogen, lower alkyl, halo-lower
 20 alkyl, lower alkoxy, halo-lower alkoxy and optionally protected
 hydroxy, and at least two of t, u, v and w are said methine group;

X is -N(SO₂R⁴)-, -N(COR⁵)- or -CO- ;

Y is -C(R⁶)(R⁷)-, -O- or -N(R⁸)-, provided that the
 compound (X) when E is nitrogen, n is 0, X is -CO- and Y is -O-
 25 is excluded) with a compound of the formula (XI):



(wherein Ar^{1p} is aryl or heteroaryl, any of which is optionally
 substituted by a substituent selected from the group consisting
 of halogen, nitro, lower alkyl, halo-lower alkyl, cyclo-lower


alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower
 alkylthio, lower alkylsulfonyl, lower alkanoyl, lower
 alkoxycarbonyl, lower alkanoylamino, $-Q^{2p}-Ar^{2p}$, optionally
 protected hydroxy, optionally protected hydroxy-lower alkyl
 5 and optionally protected carboxyl;

Ar^{2p} is aryl or heteroaryl, any of which is optionally
 substituted by a substituent selected from the group consisting
 of halogen, cyano, lower alkyl, halo-lower alkyl, lower alkoxy,
 halo-lower alkoxy, lower alkylamino, di-lower alkylamino,
 10 lower alkanoyl, aryl, optionally protected hydroxy-lower alkyl
 and optionally protected hydroxy;

Q^{2p} is a single bond, oxygen, optionally protected
 carbonyl or $-N(R^3)-$;

R^3 is hydrogen or lower alkyl;

15 Met is a common organometallic atom) in the presence of
 a catalyst,
 and optionally removing the protecting group(s) from the
 resulting compound of the formula (XII):




20 (wherein a^1 , b^1 , c^1 and d^1 are independently methine or nitrogen,
 said methine being optionally substituted by a substituent
 selected from the group consisting of halogen, cyano, lower
 alkyl, halo-lower alkyl, lower alkoxy, halo-lower alkoxy, lower
 alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy,
 25 $-N(R^{1p})R^{2p}$, $-Ar^{1p}$ and optionally protected hydroxy, and at least

one of a^1 , b^1 , c^1 and d^1 is said methine having a group represented by $-Ar^{1p}$;

Ar^{1p} , E^p , n , P , R^{1p} , R^{2p} , t , u , v , w , X and Y have each the same meaning as defined above) and/or oxidizing nitrogen atom(s) of the compound (XII).

29. A neuropeptide Y receptor antagonist agent comprising a compound of the formula (I):



(wherein A , B , C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$, and at least one of A , B , C and D is said methine group;

Ar^1 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxy carbonyl, lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting

of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;

5 n is 0 or 1;

Q^1 and Q^2 are independently a single bond, oxygen, carbonyl or $-N(R^3)-$;

R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be
10 intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;


15 R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;

T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least
20 two of T, U, V and W are said methine group;

X is $-N(SO_2R^4)-$, $-N(COR^5)-$ or $-CO-$;

Y is $-C(R^6)(R^7)-$, $-O-$ or $-N(R^8)-$, provided that the compound (I) when E is nitrogen, n is 0, X is $-CO-$ and Y is $-O-$
25 is excluded), or a salt, ester or N-oxide derivative thereof as an active ingredient.

30. An agent for treatment of bulimia, obesity or diabetes, comprising a compound of the formula (I):



(wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent
 5 selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, -N(R¹)R² and -Q¹-Ar¹, and at least one of A, B, C and D is said methine group;

10 Ar¹ is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower
 15 alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino and -Q²-Ar²;

Ar² is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower
 20 alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;

n is 0 or 1;

Q¹ and Q² are independently a single bond, oxygen, carbonyl
 25 or -N(R³)-;

R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

5 R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;

R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;


10 T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group;

15 X is $-N(SO_2R^4)-$, $-N(COR^5)-$ or $-CO-$;

Y is $-C(R^6)(R^7)-$, $-O-$ or $-N(R^8)-$, provided that the compound (I) when E is nitrogen, n is 0, X is $-CO-$ and Y is $-O-$ is excluded), or a salt, ester or N-oxide derivative thereof as an active ingredient.

20

31. A method for treatment of bulimia, obesity or diabetes, which comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of the formula (I):



25

(wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower
5 alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxycarbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-N(R^1)R^2$ and $-Q^1-Ar^1$, and at least one of A, B, C and D is said methine group;

Ar^1 is aryl or heteroaryl, any of which is optionally
10 substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl,
15 lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower
20 alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;
n is 0 or 1;

Q^1 and Q^2 are independently a single bond, oxygen, carbonyl or $-N(R^3)-$;

25 R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^8 are independently hydrogen, lower alkyl, aralkyl or aryl;


R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;

5 T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group;

10 X is $-\text{N}(\text{SO}_2\text{R}^4)-$, $-\text{N}(\text{COR}^5)-$ or $-\text{CO}-$;

Y is $-\text{C}(\text{R}^6)(\text{R}^7)-$, $-\text{O}-$ or $-\text{N}(\text{R}^8)-$, provided that the compound (I) when E is nitrogen, n is 0, X is $-\text{CO}-$ and Y is $-\text{O}-$ is excluded), or a salt, ester or N-oxide derivative thereof.

15 32. Use of a compound of the formula (I):



(wherein A, B, C and D are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkoxy carbonyl, lower alkylsulfonyl, lower alkylsulfonyloxy, $-\text{N}(\text{R}^1)\text{R}^2$ and $-\text{Q}^1-\text{Ar}^1$, and at least one of A, B, C and D is said methine group;

25 Ar^1 is aryl or heteroaryl, any of which is optionally

substituted by a substituent selected from the group consisting of halogen, nitro, hydroxy, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, cyclo-lower alkyl, lower alkenyl, lower alkoxy, halo-lower alkoxy, lower alkylthio, lower alkylsulfonyl, carboxyl, lower alkanoyl, lower alkoxycarbonyl, lower alkanoylamino and $-Q^2-Ar^2$;

Ar^2 is aryl or heteroaryl, any of which is optionally substituted by a substituent selected from the group consisting of halogen, cyano, lower alkyl, halo-lower alkyl, hydroxy-lower alkyl, hydroxy, lower alkoxy, halo-lower alkoxy, lower alkylamino, di-lower alkylamino, lower alkanoyl and aryl;

E is nitrogen, methine or hydroxy substituted methine;

n is 0 or 1;

Q^1 and Q^2 are independently a single bond, oxygen, carbonyl or $-N(R^3)-$;

R^1 and R^2 are independently hydrogen or lower alkyl, or R^1 and R^2 , taken together, form lower alkylene which may be intervened by oxygen, sulfur or imino;

R^3 is hydrogen or lower alkyl;

R^4 is lower alkyl, aralkyl or aryl;

R^5 and R^6 are independently hydrogen, lower alkyl, aralkyl or aryl;

R^6 and R^7 are independently hydrogen, hydroxy, lower alkyl, aralkyl or aryl;

T, U, V and W are independently methine or nitrogen, said methine being optionally substituted by a substituent selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, hydroxy, lower alkoxy and halo-lower alkoxy, and at least two of T, U, V and W are said methine group;

X is $-\text{N}(\text{SO}_2\text{R}^4)-$, $-\text{N}(\text{COR}^5)-$ or $-\text{CO}-$;

Y is $-\text{C}(\text{R}^6)(\text{R}^7)-$, $-\text{O}-$ or $-\text{N}(\text{R}^8)-$, provided that the compound (I) when E is nitrogen, n is 0, X is $-\text{CO}-$ and Y is $-\text{O}-$ is excluded), or a salt, ester or N-oxide derivative thereof for manufacturing
5 a medicament for treatment of bulimia, obesity or diabetes.